

Locally Non-spherical Artin Groups

Andrew Chermak

Department of Mathematics, Kansas State University, Manhattan, Kansas 66506

Communicated by I. Tite

metadata, citation and similar papers at core.ac.uk

INTRODUCTION

Let S be a non-empty set and let M be an $S \times S$ matrix (i.e., a set indexed by $S \times S$) with entries in $\mathbb{N} \cup \{\infty\}$. We write $M = (m_{a,b})_{S \times S}$. We will say that M is a *Coxeter matrix* if M is symmetric, with $m_{a,b} \geq 2$ if and only if $a \neq b$.

Let $F = F(S)$ be the free group with free basis S . For any $m \in \mathbb{N}$ and any $a, b \in S$, let $w(a, b; m)$ denote the element of F which consists of the alternating product of the generators a and b , starting with a , and having length m . Put $w(a, b; \infty) = 1$, the identity element of F . If M is a Coxeter matrix over S then the *Artin group* $G = G(M)$ is the group presented by S subject to the relations

$$w(a, b; m_{a,b}) = w(b, a; m_{a,b})$$

for all $a, b \in S$. We say further that M and G are *locally non-spherical* if for each three-element subset $\{a, b, c\}$ of S we have

$$\frac{1}{m_{a,b}} + \frac{1}{m_{a,c}} + \frac{1}{m_{b,c}} \leq 1. \quad (0.1)$$

An equivalent way to formulate this condition is to say that for any three distinct elements a, b, c of S , the Coxeter group

$$\langle a, b, c | a^2 = b^2 = c^2 = (ab)^{m_{a,b}} = (ac)^{m_{a,c}} = (bc)^{m_{b,c}} = 1 \rangle$$

is infinite.

Our aim in this paper is to prove the following two results.

THEOREM 1. *Let $G(M)$ be a locally non-spherical Artin group, where M is a Coxeter matrix over a finite set S . Then $G(M)$ has a solvable Word Problem.*

THEOREM 2. *Let $G = G(M)$ be a locally non-spherical Artin group, M a Coxeter matrix over S . Let T be a non-empty subset of S , let M_T be the submatrix of M indexed by $T \times T$, and let G_T denote the subgroup of G generated by the canonical image of T in G . Then G_T is canonically isomorphic to $G(M_T)$.*

An algorithm \mathcal{A} for the Word Problem in $G(M)$ is given explicitly in (6.5), below. It is closely related to Dehn's algorithm in that it never increases the length of a word.

The hypothesis that G be locally non-spherical generalizes the hypothesis, made in [2, 1] that G be of "large type," i.e., that $m_{a,b}$ be strictly greater than 2 for all $a, b \in S$ with $a \neq b$. Here we permit $m_{a,b} = 2$ (and we even also permit $m_{b,c} = 2$, with $c \neq a$, provided that $m_{a,c} = \infty$). On the other hand, our results are appreciably weaker than those of [2, 1]. In particular, our results do not suffice to establish solvability of the Conjugacy Problem, nor are we able to show that $G(M)$ is torsion-free.

The method employed here is an "arboreal small-cancellation theory" that was first introduced in [4] and which has undergone some improvement in [5]. This means that we study a set \mathcal{H} of hyperbolic isometries of a tree X , with $\mathcal{H} = \mathcal{H}^{-1}$, and with $\mathcal{H} \triangleleft F$ for a suitable subgroup F of $\text{Isom}(X)$. Here F will be the free group $\overline{F(S)}$, but viewed as a free product of infinite cyclic groups

$$F = \ast_{a \in S} \langle a \rangle.$$

We then take X to be the "standard tree" for this free product (and which is quite different from the Cayley graph of F with respect to S). For any $a, b \in S$ with $a \neq b$, put

$$h_{a,b} = w(a, b; m_{a,b})w(b, a; m_{a,b})^{-1}. \quad (0.2)$$

Let $N_{a,b}$ denote the normal closure of $h_{a,b}$ in the subgroup $\langle a, b \rangle$ of F generated by a and b . Then $\langle a, b \rangle / N_{a,b}$ is a two-generator Artin group, for which plenty of information is available in [2, 1], and more recently from [3]. Next, for any a, b as above, we set

$$\mathcal{H}_{a,b} = \bigcup \{g^{-1}N_{a,b}g : g \in F\} - \{1\},$$

and then define

$$\mathcal{H} = \bigcup \{\mathcal{H}_{a,b} : a, b \in S, a \neq b\}.$$

Finally, set

$$N = \langle \mathcal{H} \rangle, \quad G = F/N. \quad (0.3)$$

Then G may be identified with the Artin group $G(M)$.

Now in fact \mathcal{H} is a set of hyperbolic isometries of X . That is, for any $h \in \mathcal{H}$, h fixes no vertex and inverts no edge of X . For any such h we define

$$a(h) = \min\{d(x, x \cdot h)\}_{x \in X},$$

$$A(h) = \{x \in X: d(x, x \cdot h) = a(h)\}.$$

We call $a(h)$ the *amplitude* of h , or the *hyperbolic length* of h , and we call $A(h)$ the *axis* of h . The following two quoted results (which are in fact quite elementary) are fundamental to what follows.

0.4. PROPOSITION [7; 6, Proposition 24, p. 63]. *Let h be a hyperbolic isometry of a tree X . Then:*

- (a) $A(h)$ is a regular subtree of X with valency equal to 2.
- (b) h induces a translation of $A(h)$ of amplitude $a(h)$.
- (c) Every $\langle h \rangle$ -invariant subtree of X contains $A(h)$.
- (d) If a vertex x of X is at distance d from $A(h)$ then $d(x, xh) = a(h) + 2d$ (see Fig. 1).

0.5. PROPOSITION [6, Proposition 25, p. 63]. *Let g be an isometry of a tree X . Then the following conditions are equivalent.*





FIGURE 1

(i) g is hyperbolic.

(ii) There is an edge e of X such that g translates e a positive distance along a geodesic path of X .



Moreover, if (ii) holds for the edge e then e is an edge of $A(g)$.

The bulk of the present paper is devoted to showing that \mathcal{H} has a certain kind of “convergence property” (as defined below, in (2.5)). For a discussion of “convergence” see the introduction to [4] and see especially [5]. This will occupy Sections 2 through 5. Section 6 is then a straightforward translation of this convergence property into Theorems 1 and 2.

Section 2 contains a few technical results concerning operations on finite sequences of isometries, as well as a review of some definitions from [5]. In particular, we remind the reader of the notion of a “polarization” of X over \mathcal{H} in (2.6), and of what it means for a polarization \mathcal{P} to “control convergence” in (2.8). After this, Section 3 is then possibly the most difficult to read, since it consists mostly of a chain of intricate definitions, culminating in the definition (3.12) of a suitable polarization \mathcal{P} . Sections 4 and 5 then perform the task of verifying that \mathcal{P} controls convergence.

1. PRELIMINARIES

Fix a Coxeter matrix $M = (m_{a,b})_{S \times S}$, and let F denote the free group with free basis S . Define a graph X as follows. The vertex set $V(X)$ will be the (disjoint) union of two sets D and D' where D is simply the set F of group elements, and where D' is the set of all right cosets $\langle a \rangle g$ for $a \in S$ and $g \in F$. Then define an incidence relation between D and D' by means of membership: the element h of D and the element $\langle a \rangle g$ of D' are incident if $h \in \langle a \rangle g$, i.e., $h = a^n g$ for some n . Thus, X is a bipartite graph. In fact, Bass–Serre theory [6] shows that X is a tree, the “standard tree” for the free product F of the cyclic groups $\langle a \rangle$, $a \in S$.

Let d be the natural metric on $V(X)$. It will be convenient to identify X with its topological realization, so that X becomes a simplicial tree, with edges of length 1. Vertices of X will also be called *points*. For any points x and y in X , $[x, y]$ denotes the oriented closed path from x to y in X . We call this the *oriented closed segment* from x to y . If $J = [x, y]$ is an oriented closed segment we put

$$\partial_0 J = \{x\}, \quad \partial_1 J = \{y\}, \quad \partial J = \{x, y\}.$$

We may regard $[x, y]$ as a certain subset of X , together with a preferred direction. The underlying subset of $[x, y]$ is called the *closed segment* joining x and y , and it too will be denoted $[x, y]$. No confusion need result from this abuse of notation.

A closed segment is *degenerate* if it consists of a single point. We write $[x, y]$ for the subset $[x, y] - \{y\}$ of X , and we similarly define $(x, y]$ and (x, y) . If $J = [x, y]$ is a closed segment, we write $l(J)$ for the *length* of J , which is $d(x, y)$.

Notice that F acts on X by right translation, and in this way every element of F may be viewed as an isometry of X .

1.1. LEMMA. (a) *Edge-stabilizers in F are trivial. Indeed, the stabilizer in F of any vertex in D is trivial.*

(b) *F is a subgroup of $\text{Isom}(X)$.*

Proof. Part (a) is immediate, and (b) follows from (a).

Let $1 \neq g \in F$. We can then write

$$g = a_{i_1}^{e_1} \cdots a_{i_m}^{e_m},$$

$$\text{with } a_{i_j} \in S, e_j \in \mathbb{Z} - \{0\}, \text{ and with } a_{i_j} \neq a_{i_{j+1}} \text{ for any } j. \quad (1.2)$$

We call (1.2) the *normal form* of g , and we say that g is *cyclically reduced* if also $a_{i_m} \neq a_{i_1}$. Now set

$$|g| = |e_1| + \cdots + |e_m|, \quad \text{and} \quad \|g\| = m.$$

We refer to $|g|$ as the *length* of g , and to $\|g\|$ as the *syllable-length* of g .

1.3. LEMMA. *Let $g \in F$, with normal form as in (1.2). Then the vertex-sequence for the oriented closed segment $[1, g]$ is*

$$(1, \langle a_{i_m} \rangle, a_{i_m}^{e_m}, \langle a_{i_{m-1}} \rangle a_{i_m}^{e_m}, \dots, \langle a_{i_1} \rangle a_{i_2}^{e_2} \cdots a_{i_m}^{e_m}, g).$$

In particular, we have $d(1, g) = 2 \cdot \|g\|$.

Proof. This is immediate.

For any $a, b \in S$ with $a \neq b$, we have defined (0.2) the basic relator:

$$h_{a,b} = w(a, b; m_{a,b}) w(b, a; m_{a,b})^{-1}.$$

Denote by $N_{a,b}$ the smallest normal subgroup of $\langle a, b \rangle$ containing $h_{a,b}$. Thus, $\langle a, b \rangle / N_{a,b}$ is the Artin group with matrix

$$\begin{pmatrix} 1 & m_{a,b} \\ m_{a,b} & 1 \end{pmatrix}.$$

We require the following important result from [2].

1.4 [2, Lemma 7, p. 210]. *Let $g_1, g_2 \in \langle a, b \rangle$, and suppose that $g_1 g_2 \in N_{a,b}$.*

- (a) *If $\|g_1\| \leq m_{a,b}$ then $|g_1| \leq |g_2|$.*
- (b) *If $\|g_1\| < m_{a,b}$ then $|g_1| < |g_2|$ or $g_1 = g_2^{-1}$.*

1.5. COROLLARY (which is also [2, Lemma 6]). *For any $g \in N_{a,b}$, $g \neq 1$, we have $\|g\| \geq 2 \cdot m_{a,b}$.*

Let $X_{a,b}$ denote the subtree of X induced on the vertex set $\{g, \langle a \rangle g, \langle b \rangle g : g \in \langle a, b \rangle\}$. Certainly, $X_{a,b}$ is invariant under $\langle a, b \rangle$. Let h be a cyclically reduced element of $N_{a,b}$, $h \neq 1$. After possibly reversing the roles of a and b , we may then write $h = a^{e_1} b^{f_1} \dots a^{e_l} b^{f_l}$, for some non-zero integers e_i and f_i , $1 \leq i \leq l$. By (1.3), the vertex-sequence for the oriented closed segment $[1, h]$ is then

$$(1, \langle b \rangle, b^{f_1}, \langle a \rangle b^{f_1}, \dots, \langle a \rangle b^{f_1} \dots a^{e_l} b^{f_l}, h),$$

and $d(1, h) = 2\|h\| = 4l$. Observe that $\langle a \rangle h = \langle a \rangle b^{f_1} \dots a^{e_l} b^{f_l}$, so that the closed segment $[\langle a \rangle, 1]$ is translated a distance $2 \cdot \|h\|$ by h , along the segment $[1, h]$. Now (0.5) yields the following result.

1.6. LEMMA. *Let h be a cyclically reduced element of $N_{a,b}$, $h \neq 1$. Then h is a hyperbolic isometry of X , and $a(h) = 2 \cdot \|h\|$. Further, we have $A(h) \subseteq X_{a,b}$, and $A(h)$ contains the closed segment $[\langle a \rangle, \langle b \rangle]$ (whose vertex-sequence is $(\langle a \rangle, 1, \langle b \rangle)$).*

For any $a, b \in S$ with $a \neq b$, put

$$\mathcal{H}_{a,b} = \bigcup_{g \in F} \{g^{-1} N_{a,b} g\} - \{1\},$$

and then put

$$\mathcal{H} = \bigcup \{\mathcal{H}_{a,b} : a, b \in S, a \neq b\}.$$

Let $h \in \mathcal{H}$. Then h is conjugate to a cyclically reduced element of $N_{a,b}$, for some a and b , and hence h is a hyperbolic isometry of X , by (1.6). Now let g be a vertex in $D \cap A(h)$. The two neighbors of g in $A(h)$ are then

vertices $\langle c \rangle g$ and $\langle d \rangle g$ for some $c, d \in S$ with $c \neq d$. Then $[\langle c \rangle, \langle d \rangle]$ is a segment of $A(h^{s^{-1}})$. On the other hand, we are given that there is a conjugate $h' = h^{s'}$ of h in $N_{a,b}$. Without loss, h' is cyclically reduced, and then $[\langle a \rangle, \langle b \rangle]$ is a segment of $A(h')$. Now gg' maps $A(h^{s^{-1}})$ onto $A(h')$, so gg' maps 1 into $X_{a,b}$. Thus, $gg' \in \langle a, b \rangle$, and so $\{c, d\} = \{a, b\}$. This yields parts (a) and (b) of the following result.

1.7. LEMMA. (a) \mathcal{H} is the disjoint union of the sets $\mathcal{H}_{a,b}$, $a, b \in S$, $a \neq b$.

(b) Let $h \in \mathcal{H}$. Then $h \in \mathcal{H}_{a,b}$ if and only if there exists $g \in F$ such that $[\langle a \rangle, \langle b \rangle] \cdot g$ is a closed segment of $A(h)$.

(c) Let $h \in \mathcal{H}_{a,b}$ and let $c \in S$. Suppose that $\langle c \rangle$ is a vertex of $A(h)$. Then $c \in \{a, b\}$ and $h \in N_{a,b}$.

(d) If $a, b \in S$ with $a \neq b$, and $[\langle a \rangle, \langle b \rangle]$ is a closed segment of $A(h)$, where $h \in \mathcal{H}$, then $h \in N_{a,b}$.

Proof. We need only prove (c) and (d), and we note that (d) is an immediate consequence of (c). Now let $h \in \mathcal{H}_{a,b}$. It then follows from (a) and (b) that every vertex of $A(h)$ which is not in D is of the form $\langle a \rangle g$ or $\langle b \rangle g$ for some $g \in F$. If $c \in S$ and $\langle c \rangle$ is a vertex of $A(h)$, it now follows that $c \in \{a, b\}$. Without loss, $c = a$. Now $\langle a \rangle$ acts transitively on the vertices adjacent to the vertex $\langle a \rangle$, and since $N_{a,b}$ is $\langle a \rangle$ -invariant we may then assume (after possibly replacing h by h^{a^n} for some n) that 1 is a vertex of $A(h)$. By assumption, $h^g \in N_{a,b}$ for some $g \in F$. Then g maps 1 into $X_{a,b}$, so $g \in \langle a, b \rangle$ and $h \in N_{a,b}$, proving (c).

We now come to two “Small Cancellation” results, which place restrictions on the ways in which axes for elements of \mathcal{H} can intersect.

1.8. LEMMA. Let $h, h' \in \mathcal{H}$ with $hh' \notin \mathcal{H} \cup \{1\}$. Then $l(A(h) \cap A(h')) \leq 2$, with equality only if $\partial(A(h) \cap A(h')) \subseteq D$.

Proof. Suppose this is false. Then $A(h) \cap A(h')$ contains a closed segment I of length 2, with $\partial I \subseteq D'$. By (1.7)(b) we may replace h and h' by h^g and $(h')^g$ for a suitable $g \in F$, obtaining $I = [\langle a \rangle, \langle b \rangle]$ for some $a, b \in S$. But then $hh' \in N_{a,b}$ by (1.7)(d), and so $hh' \in \mathcal{H} \cup \{1\}$, for a contradiction.

In order to state the next result, we need to introduce the following notation. For any $h \in \mathcal{H}$, we put

$$m(h) = m_{a,b},$$

where $\{a, b\}$ is the uniquely determined subset of S such that $h \in \mathcal{H}_{a,b}$.

(1.9)

We will now make use of the hypothesis that our Artin group G is locally non-spherical.

1.10. LEMMA. *Let h_1, h_2, h_3 be three distinct elements of \mathcal{H} . Suppose that for all i and j with $1 \leq i < j \leq 3$, we have*

(i) $h_i h_j \notin \mathcal{H} \cup \{1\}$, and

(ii) $A(h_i) \cap A(h_j)$ is a non-degenerate closed segment, oriented oppositely by h_i and h_j .

Then $1/m(h_1) + 1/m(h_2) + 1/m(h_3) \leq 1$.

Proof. The conditions guarantee that there are closed segments $[x, y]$ of $A(h_1)$, $[y, z]$ of $A(h_2)$, and $[z, x]$ of $A(h_3)$, all of length 2, arranged as in Fig. 2.

Let w denote the vertex at the center of the picture. That is, $\{w\} = A(h_1) \cap A(h_2) \cap A(h_3)$. If we conjugate all three elements h_1, h_2 , and h_3 by an element g of F , then the condition (i) remains invariant, as do the integers $m(h_i)$. We may therefore assume that either $w = 1$ or $w = \langle a \rangle$ for some $a \in S$.

Suppose $w = 1$. Then there are distinct elements a, b, c of S with $x = \langle a \rangle$, $y = \langle b \rangle$, and $z = \langle c \rangle$. By (1.7)(d) we get $h_1 \in N_{a,b}$, $h_2 \in N_{b,c}$, and $h_3 \in N_{a,c}$. As G is locally non-spherical, we have

$$\frac{1}{m_{a,b}} + \frac{1}{m_{a,c}} + \frac{1}{m_{b,c}} \leq 1$$

as desired.

Suppose $w = \langle a \rangle$ for some $a \in S$. By (1.7)(c) we then have $h_1 \in N_{a,b}$, $h_2 \in N_{a,c}$, and $h_3 \in N_{a,d}$, for some elements b, c, d of S . As $h_i h_j \notin \mathcal{H} \cup \{1\}$ for any $i \neq j$, it follows that b, c , and d are pairwise distinct. This then yields

$$\frac{1}{m_{a,b}} + \frac{1}{m_{a,c}} + \frac{1}{m_{a,d}} \leq 1,$$

as required.




FIGURE 2

1.11. For any $h \in H$, put $b(h) = 4 \cdot m(h)$. Also, fix an isometry α_h from \mathbb{Z} to $A(h)$, such that α_h carries the even integers onto $D \cap A(h)$. (Here we view \mathbb{Z} as a simplicial tree, with edges $\{n, n+1\}$ of length 1.) The map \mathcal{D} which assigns to each $h \in \mathcal{H}$ the triple $(\alpha_h, 2\mathbb{Z}, D \cap A(h))$ is then a *specialization* on \mathcal{H} , in the sense of [5, Definition (3.7)].

Observe that $b(h) \in 4\mathbb{Z}$, and $b(h) = b(h^{-1}) = b(h^g)$ for any $g \in F$. If we can show also that $b(h) \leq a(h)$, then $b(h)$ will be a *modified length function* on \mathcal{H} , in the sense of [5, Definition (3.8)]. In fact, such is the case, as we now prove.

1.12. LEMMA. *Let $h \in \mathcal{H}$. Then $a(h) = 2 \cdot \|h'\|$ for any cyclically reduced conjugate h' of h , and $a(h) \geq b(h)$. In particular, $a(h) \geq 8$.*

Proof. Since $a(h) = a(h')$ and $b(h) = b(h')$ for any conjugate h' of h , we may assume that h is cyclically reduced, and that $h \in N_{a,b}$ for some $a, b \in S$, $a \neq b$. Then $a(h) = 2\|h\|$ by (1.6), and so (1.5) yields $a(h) \geq 4 \cdot m_{a,b} = b(h)$.

2. \mathcal{H} -SEQUENCES

As in [5], an \mathcal{H} -sequence is always a finite sequence $\mathbf{h} = (h_0, \dots, h_n)$ of elements of \mathcal{H} . We write $\mathbf{h}^{-1} = (h_n^{-1}, \dots, h_0^{-1})$, and for any $g \in \langle \mathcal{H} \rangle$ we write \mathbf{h}^g for (h_0^g, \dots, h_n^g) . (Here x^g denotes $g^{-1}xg$ for any $x \in \mathcal{H}$.) We write $l(\mathbf{h})$ for the *length* of \mathbf{h} , which is here $n+1$.

If \mathbf{h} and \mathbf{h}' are \mathcal{H} -sequences then $\mathbf{h} \circ \mathbf{h}'$ denotes the \mathcal{H} -sequence obtained by concatenating \mathbf{h} and \mathbf{h}' . A *segment* of \mathbf{h} is an \mathcal{H} -sequence \mathbf{s} such that $\mathbf{h} = \mathbf{x} \circ \mathbf{s} \circ \mathbf{y}$ for some \mathcal{H} -sequences \mathbf{x} and \mathbf{y} . If $\mathbf{x} = (\emptyset)$ we say that \mathbf{s} is a *prefix* of \mathbf{h} , and we say that \mathbf{s} is a *suffix* of \mathbf{h} if $\mathbf{y} = (\emptyset)$.

The \mathcal{H} -sequence \mathbf{h} is *reduced* if for all i , $1 \leq i \leq n$, we have $h_{i-1}h_i \notin \mathcal{H} \cup \{1\}$. We put $\mathcal{H}^{**} = \{\mathbf{h} : \mathbf{h} \text{ is a reduced } \mathcal{H}\text{-sequence}\}$. We recall the notion of *braiding* of \mathcal{H} -sequences, from Section 3 of [5]. Thus, the \mathcal{H} -sequence $\mathbf{h}' = (h'_0, \dots, h'_n)$ is a *simple braiding* of \mathbf{h} if there exists k with $1 \leq k \leq n$ such that $h'_i = h_i$ for all $i \notin \{k-1, k\}$, and either

$$(h'_{k-1}, h'_k) = (h_{k-1}h_k h_{k-1}^{-1}, h_{k-1}),$$

or

$$(h'_{k-1}, h'_k) = (h_k, h_k^{-1}h_{k-1}h_k).$$

If there exists a chain of \mathcal{H} -sequences

$$\mathbf{h} = \mathbf{s}_0, \dots, \mathbf{s}_l = \mathbf{h}'$$

such that \mathbf{s}_i is a simple braiding of \mathbf{s}_{i-1} for all i , $1 \leq i \leq l$, then \mathbf{h}' is a braiding of \mathbf{h} . In fact, there is an action of the Artin braid group B_{n+1} (on $n+1$ strings) on the set of \mathcal{H} -sequences of length $n+1$, such that the n standard generators of B_{n+1} , and their inverses, perform the simple braidings. We see then that \mathbf{h}' is a braiding of \mathbf{h} if and only if \mathbf{h}' lies in the same orbit as \mathbf{h} under the action of B_{n+1} . We put

$$\mathcal{H}^* = \{\mathbf{h} : \mathbf{h} \text{ is an } \mathcal{H}\text{-sequence, and every braiding of } \mathbf{h} \text{ is in } \mathcal{H}^{**}\}.$$

We quote the following result from [5, Lemma (3.6)].

2.0. LEMMA. (a) Let $\mathbf{h} = (h_0, \dots, h_n)$ be a word of minimal length, in the alphabet \mathcal{H} , for the element $h_0 \cdots h_n$ of $\text{Isom}(X)$. Then $\mathbf{h} \in \mathcal{H}^*$.

(b) Let $\mathbf{h} = (h_0, \dots, h_n)$ and $\mathbf{h}' = (h'_0, \dots, h'_n)$ be \mathcal{H} -sequences, and suppose that h' is a braiding of \mathbf{h} . Then $h_0 \cdots h_n = h'_0 \cdots h'_n$.

2.1. DEFINITION. Let \mathbf{h} and \mathbf{h}' be \mathcal{H} -sequences, with $\mathbf{h} = (h_0, \dots, h_n)$, and let m be an integer, $0 \leq m \leq n$. We say that \mathbf{h}' is a *simple distillation* of \mathbf{h} (of degree m) if there exist k and l with $0 \leq k \leq l \leq m$, such that

$$\mathbf{h}' = (h_0, \dots, h_{k-1})^{h_m} \circ (h_m) \circ (h_l, \dots, h_{m-1}) \circ (h_{m+1}, \dots, h_n).$$

In words: \mathbf{h}' is obtained from \mathbf{h} by permuting h_m to the left, across (h_k, \dots, h_{m-1}) , deleting (h_k, \dots, h_{l-1}) , and conjugating (h_0, \dots, h_{k-1}) by h_m . Notice that

$$l(h') = (n+1) - (l-k).$$

We will also say that the *codegree* of the above simple distillation is $l(\mathbf{h}) - m$.

EXAMPLE. Some simple distillations of (h_0, h_1, h_2, h_3) of degree 2 (and codegree 2) are (h_2, h_3) , $(h_0^{h_2}, h_2, h_3)$, $(h_0^{h_2}, h_2, h_1, h_3)$, and (h_2, h_0, h_1, h_3) .

2.2. DEFINITION. Let \mathbf{h} and \mathbf{h}' be \mathcal{H} -sequences, with $\mathbf{h} = (h_0, \dots, h_n)$. We say that \mathbf{h}' is a *distillation* of \mathbf{h} if $\mathbf{h} = \mathbf{h}' = (\emptyset)$, or if there exists a chain of \mathcal{H} -sequences

$$\mathbf{h} = \mathbf{s}_0, \dots, \mathbf{s}_n = \mathbf{h}' \quad (n+1 = l(\mathbf{h})),$$

such that for all m with $1 \leq m \leq n$, \mathbf{s}_m is a simple distillation of \mathbf{s}_{m-1} of codegree $n - m + 1$.

EXAMPLE. The sequence $(h_1^{h_2 h_3}, h_3)$ is a distillation of $(h_0, h_1, h_2, h_3) = \mathbf{s}_0$, as is revealed by taking $\mathbf{s}_1 = (h_1, h_2, h_3)$, $\mathbf{s}_2 = (h_1^{h_2}, h_2, h_3)$, and $\mathbf{s}_3 = (h_1^{h_2 h_3}, h_3)$. Here $n+1 = 4$, and the reader may check, for example, that the codegree of the simple distillation from \mathbf{s}_2 to \mathbf{s}_3 is $1 = 4 - 3$.

2.3. LEMMA. *Let $\mathbf{h} \in \mathcal{H}^*$, let \mathbf{h}' be a distillation of \mathbf{h} , and let $h \in \mathcal{H}$ such that $\mathbf{h} \circ (h) \in \mathcal{H}^*$. Then $\mathbf{h}' \in \mathcal{H}^{**}$, and for any entry h' of \mathbf{h}' we have $hh' \notin \mathcal{H} \cup \{1\}$ and $h'h \notin \mathcal{H} \cup \{1\}$.*

Proof. By assumption there is a chain

$$\mathbf{h} = \mathbf{s}_0, \dots, \mathbf{s}_n = \mathbf{h}'$$

such that each \mathbf{s}_m is a simple distillation of \mathbf{s}_{m-1} of codegree $n - m + 1$, where $n + 1 = l(\mathbf{h})$. Write $\mathbf{h} = (h_0, \dots, h_n)$, as usual. Fix $m \geq 1$ and write

$$\mathbf{s}_{m-1} = (g_0, \dots, g_r), \quad \mathbf{s}_m = (g'_0, \dots, g'_s),$$

where of course $n \geq r \geq s$. Put $m' = r + m - n$. We aim to prove the following statements, by induction on m .

- (1)' $\mathbf{s}_m = (g'_0, \dots, g'_{s+m-n}) \circ (h_{m+1}, \dots, h_n)$.
- (2)' $(g'_0, \dots, g'_{s+m-n}) \in \mathcal{H}^{**}$.
- (3)' $g'_i g'_j \notin \mathcal{H} \cup \{1\}$ for any i and j with $i \neq j > s + m - n$.

Notice that if $m = 0$ then $s = n$, and then (1)' and (2)' are obviously true, while $h_i h_j \notin \mathcal{H} \cup \{1\}$ for any $i \neq j$, by [5, Lemma (3.6)(e)]. We may now assume that versions of (1)' through (3)' hold for \mathbf{s}_{m-1} . Thus, we have:

- (1) $\mathbf{s}_{m-1} = (g_0, \dots, g_{r+m-n-1}) \circ (h_m, \dots, h_n)$.
- (2) $(g_0, \dots, g_{r+m-n-1}) \in \mathcal{H}^{**}$.
- (3) $g_i g_j \notin \mathcal{H} \cup \{1\}$ for any i and j with $i \neq j > r + m - n - 1$.

Notice that $g_{m'} = h_m$. Since \mathbf{s}_m is a simple distillation of \mathbf{s}_{m-1} of codegree $n - m + 1$, there exist integers k and l with $0 \leq k \leq l \leq m'$, such that

$$\mathbf{s}_m = (g_0, \dots, g_{k-1})^{h_m} \circ (h_m) \circ (g_l, \dots, g_{m'-1}) \circ (h_{m+1}, \dots, h_n).$$

Since $s = r - l + k$, we have $s + m - n = m' - l + k$ and so (1)' holds. We have

$$(g'_0, \dots, g'_{s+m-n}) = (g_0, \dots, g_{k-1})^{h_m} \circ (h_m) \circ (g_l, \dots, g_{m'-1}).$$

For any $i < m' - 1$, (2) says that $g_i g_{i+1} \notin \mathcal{H} \cup \{1\}$. Then also $(g_i g_{i+1})^{h_m} \notin \mathcal{H} \cup \{1\}$. Further, we have $(g_{k-1})^{h_m} h_m = (g_{k-1} h_m)^{h_m} \notin \mathcal{H} \cup \{1\}$ since (3) tells us that $g_{k-1} h_m \notin \mathcal{H} \cup \{1\}$. Since $h_m g_l = (g_l h_m)^{h_m^{-1}}$, (3) also implies that $h_m g_l \notin \mathcal{H} \cup \{1\}$. This proves (2)'.

Observe that for any $i \leq s + m - n$, there exists a chain of integers $c < c_1 < \dots < c_u \leq m$ such that

$$g'_i = (h_c)^{h_{c_1} h_{c_2} \dots h_{c_u}}.$$

Now consider the chain of \mathcal{H} -sequences

$$\begin{aligned}
 \mathbf{h} &= \mathbf{h}_0 = (h_0, \dots, h_c, \dots, h_{c_1}, \dots, h_{c_u}, \dots, h_n) \\
 \mathbf{h}_1 &= (h_{c_1}) \circ (h_0, \dots, h_{c_1-1})^{h_{c_1}} \circ (h_{c_1+1}, \dots, h_n) \\
 \mathbf{h}_2 &= (h_{c_1}, h_{c_2}) \circ (h_0, \dots, h_{c_1-1})^{h_{c_1} h_{c_2}} \circ (h_{c_1+1}, \dots, h_{c_2-1})^{h_{c_2}} \\
 &\quad \circ (h_{c_2+1}, \dots, h_n) \\
 &\quad \vdots \\
 \tilde{\mathbf{h}} &= \mathbf{h}_u \\
 &= (h_{c_1}, h_{c_2}, \dots, h_{c_u}) \circ (h_0, \dots, h_{c_1-1})^{h_{c_1} \cdots h_{c_u}} \circ (h_{c_1+1}, \dots, h_{c_2-1})^{h_{c_2} \cdots h_{c_u}} \\
 &\quad \circ \cdots \circ (h_{c_{u-1}+1}, \dots, h_{c_u-1})^{h_{c_u}} \circ (h_{c_u+1}, \dots, h_n).
 \end{aligned}$$

This chain displays $\tilde{\mathbf{h}}$ as a braiding of \mathbf{h} . But $(h_c)^{h_{c_1} \cdots h_{c_u}}$ is an entry of $\tilde{\mathbf{h}}$, and also h_j is an entry of $\tilde{\mathbf{h}}$ for $j \geq m$. Then [5, Lemma (3.6)(e)] yields $(h_c)^{h_{c_1} \cdots h_{c_u}} h_j \notin \mathcal{H} \cup \{1\}$, and yields also $h_i h_j \notin \mathcal{H} \cup \{1\}$ if $m \leq i < j$. This completes the proof of (3)'.

Applying (2)' to the case $m = n$, we obtain $\mathbf{h}' \in \mathcal{H}^{**}$. Now replace \mathbf{h} with $\mathbf{h} \circ (h)$, and obtain a distillation of $\mathbf{h} \circ (h)$ via the sequence $\mathbf{s}_0 \circ (h), \dots, \mathbf{s}_n \circ (h)$, $(h) \circ \mathbf{s}_n = (h) \circ \mathbf{h}'$. Apply (3)' to $\mathbf{s}_n \circ (h)$ and obtain $h'h \notin \mathcal{H} \cup \{1\}$ for any entry h' of \mathbf{h}' . As $hh' = (h'h)^{h^{-1}}$, the proof of the lemma is complete.

2.4. LEMMA. *Let \mathbf{h} be an \mathcal{H} -sequence. Then there exists a braiding \mathbf{g} of \mathbf{h}^{-1} such that, for any distillation \mathbf{h}' of \mathbf{h} , \mathbf{h}'^{-1} is a distillation of \mathbf{g} .*

Proof. Notice first of all that if \mathbf{s} is an \mathcal{H} -sequence and \mathbf{t} is a simple braiding of \mathbf{s} , then \mathbf{t}^{-1} is a simple braiding of \mathbf{s}^{-1} . It follows that the inverse of a braiding of \mathbf{s} is a braiding of \mathbf{s}^{-1} .

Let $\mathbf{h} = (h_0, \dots, h_n)$. For any i , write \bar{h}_i for h_i^{-1} , and put $g_i = \bar{h}_i^{h_{i+1} \cdots h_n}$. (This means that $g_n = h_n^{-1}$). Put $\mathbf{g} = (g_0, \dots, g_n)$. Then \mathbf{g}^{-1} is a braiding of \mathbf{h} , as is displayed by the following sequence of intermediate braidings:

$$\begin{aligned}
 &(h_0, \dots, h_n), \\
 &(h_1, h_0^{h_1}, h_2, \dots, h_n), \\
 &(h_2, h_1^{h_2}, h_0^{h_1 h_2}, h_3, \dots, h_n), \\
 &\quad \vdots \\
 &(h_n, h_{n-1}^{h_n}, \dots, h_0^{h_1 \cdots h_n}) = \mathbf{g}^{-1}.
 \end{aligned}$$

The observation made in the first paragraph of this proof then says that \mathbf{g} is a braiding of \mathbf{h}^{-1} .

For any \mathcal{H} -sequence \mathbf{s} and for any j with $0 \leq j \leq l(\mathbf{s})$, the prefix \mathbf{x} of \mathbf{s} such that $l(\mathbf{x}) = l(\mathbf{s}) - t$ will be called the prefix of \mathbf{s} of *colength* t .

We are given a distillation \mathbf{h}' of \mathbf{h} . Thus, there exists a chain of \mathcal{H} -sequences

$$\mathbf{h} = \mathbf{s}_0, \dots, \mathbf{s}_n = \mathbf{h}'$$

such that \mathbf{s}_m is a simple distillation of \mathbf{s}_{m-1} of codegree $n - m + 1$ ($1 \leq m \leq n$). We may write

$$\mathbf{s}_{m-1} = (a_0, \dots, a_r) \circ (h_m, \dots, h_n)$$

for some $r < m$. Thus, (a_0, \dots, a_r) is the prefix of \mathbf{s}_{m-1} of colength $n - m + 1$. Our object now will be to produce a chain

$$\mathbf{g} = \mathbf{t}_0, \dots, \mathbf{t}_n = \mathbf{g}',$$

where \mathbf{t}_m is a simple distillation of \mathbf{t}_{m-1} of codegree $n - m + 1$ ($1 \leq m \leq n$), and where $\mathbf{g}' = \mathbf{h}'^{-1}$. We claim that there exists such a chain, with the property that for every m with $0 \leq m \leq n$, the prefix of \mathbf{t}_m of colength $n - m$ is $(\mathbf{x}^{-1})^{h_{m+1} \cdots h_n}$, where \mathbf{x} is the prefix of \mathbf{s}_m of colength $n - m$. For $m = n$, this simply says that $\mathbf{g}' = \mathbf{h}'^{-1}$, as desired. For $m = 0$, it says that $\mathbf{g}_0 = h_0^{h_1 \cdots h_n}$, which is true by definition. Proceeding by induction, we may then assume that we have

$$\begin{aligned} \mathbf{t}_{m-1} &= (b_0, \dots, b_r) \circ (g_m, \dots, g_n), \\ (b_0, \dots, b_r) &= (\bar{a}_r, \dots, \bar{a}_0)^{h_m \cdots h_n}. \end{aligned}$$

Since \mathbf{s}_m is a simple distillation of \mathbf{s}_{m-1} of codegree $n - m + 1$, we have

$$\mathbf{s}_m = (a_0, \dots, a_{k-1})^{h_m} \circ (h_m) \circ (a_l, \dots, a_r) \circ (h_{m+1}, \dots, h_n)$$

for some k and l , with $0 \leq k \leq l \leq r + 1$. Put $k' = r - l + 1$ and $l' = r - k + 1$. Then $0 \leq k' \leq l' \leq r + 1$, and we then define

$$\mathbf{t}_m = (b_0, \dots, b_{k'-1})^{g_m} \circ (g_m) \circ (b_{l'}, \dots, b_r) \circ (g_{m+1}, \dots, g_n).$$

We need to show

$$\begin{aligned} &(b_0, \dots, b_{k'-1})^{g_m} \circ (g_m) \circ (b_{l'}, \dots, b_r) \\ &= \left((\bar{a}_r, \dots, \bar{a}_l) \circ (\bar{h}_m) \circ (\bar{a}_{k-1}, \dots, \bar{a}_0)^{h_m} \right)^{h_{m+1} \cdots h_n}. \end{aligned}$$

It will be left to the reader to perform this straightforward exercise, and to thereby complete the proof of (2.4).

We end this section by reviewing some of the terminology from [5], for the convenience of the reader.

Recall the definitions of b and \mathcal{D} from (1.11).

2.5. DEFINITION. Let $\mathbf{h} = (h_0, \dots, h_n) \in \mathcal{H}^*$ and let $x \in X$. Put $x_0 = x$, and define $x_{k+1} = x_k \cdot h_k$, $0 \leq k \leq n$. Write z for x_{n+1} . Also, put $I_k = A(h_k) \cap [x_k, x_{k+1}] \cap [x_k, z]$. We say that x converges to z via h (relative to (b, \mathcal{D})) if for all k , $0 \leq k \leq n$, we have

$$d(h_{k+1}, z) + a(h_k) \leq d(x_k, z) + b(h_k),$$

with strict inequality if $\partial I_k \not\subseteq D$. (C_k)

We say that \mathcal{H} has the (b, \mathcal{D}) -convergence property if, whenever $\mathbf{h} \in \mathcal{H}^*$ and $x \in X$, there exists a braiding \mathbf{h}' of \mathbf{h} such that x converges to z via \mathbf{h}' . (Note that z is well-defined, by (2.0)(b).)

The main result of this paper, from which all else will follow, is that \mathcal{H} has the (b, \mathcal{D}) -convergence property. The proof of this will be achieved in Section 5, below.

2.6. DEFINITION. Let \mathcal{P} be a mapping which associates to each $\mathbf{h} \in \mathcal{H}^*$ a collection $\mathcal{P}(\mathbf{h})$ of closed oriented segments of X . We say that \mathcal{P} is a polarization (of X over \mathcal{H}) if the following conditions hold.

- (a) $\mathcal{P}((\emptyset))$ is the set of all degenerate closed segments of X . That is, $\mathcal{P}((\emptyset)) = X$.
- (b) $\mathcal{P}(\mathbf{h}) = \mathcal{P}(\mathbf{h}')$ if \mathbf{h}' is a braiding of \mathbf{h} .
- (c) If $J \in \mathcal{P}(\mathbf{h})$ then $J^{opp} \in \mathcal{P}(\mathbf{h}^{-1})$, and $J \cdot g \in \mathcal{P}(\mathbf{h}^g)$ for any $g \in \langle \mathcal{H} \rangle$.

2.7. DEFINITION. Let \mathcal{P} be a fixed polarization of X over \mathcal{H} , let $\mathbf{h} \in \mathcal{H}^*$, and let $J \in \mathcal{P}(\mathbf{h})$. Then $\mathcal{E}(\mathbf{h}, J)$ denotes the set of all $h \in \mathcal{H}$ satisfying the following three conditions.

- (a) $\mathbf{h} \circ (h) \in \mathcal{H}^*$,
- (b) $J \cap A(h) \neq \emptyset$, and
- (c) if J is non-degenerate, then $J \cap A(h)$ is non-degenerate, and is oriented toward $\partial_0 J$ by h .

2.8. DEFINITION. Let \mathcal{P} be a polarization of X over \mathcal{H} , and let b and \mathcal{D} be defined as in (1.11). We say that \mathcal{P} controls convergence (relative to (b, \mathcal{D})) if the following conditions hold whenever we have $\mathbf{h} \in \mathcal{H}^*$, $J \in \mathcal{P}(\mathbf{h})$, and $h \in \mathcal{E}(\mathbf{h}, J)$.

(1) (*Exclusion*)

- (a) $J \not\subseteq A(h)$ if $\mathbf{h} \neq (\emptyset)$,
- (b) $l(J \cap A(h)) < a(h)$, and
- (c) if $\partial J \cap A(h) \neq \emptyset$ then $b(h) > 2 \cdot l(J \cap A(h))$.

(2) (*Extension*)

- (a) We have $[(\partial_0 J) \cdot h, \partial_1 J] \in \mathcal{P}(\mathbf{h} \circ (h))$, provided that:
 - (i) $\partial J \cap A(h) = \emptyset$, and
 - (ii) $l(J \cap A(h)) \geq a(h) - \frac{1}{2}b(h)$, with strict inequality if $\partial(J \cap A(h)) \not\subseteq D$.
- (b) For $x \in A(h)$, we have $[x \cdot h, \partial_1 J] \in \mathcal{P}(\mathbf{h} \circ (h))$, provided that:
 - (i) $\partial_0 J \in [x, x \cdot h)$, and
 - (ii) $d(\partial_0 J, x \cdot h) \geq \frac{1}{2}b(h)$, with strict inequality if $\{\partial_0 J, x\} \subseteq D$.

3. A POLARIZATION OF X

Let $J = [x, x']$ be a non-degenerate, oriented closed segment of X . Recall from [5, Sect. 2] that this means that J consists of a metric morphism $J: [0, d(x, x')] \rightarrow X$ with $0 \cdot J = x$ and with $d(x, x') \cdot J = x'$. We also write $\partial_0 J$ for x and $\partial_1 J$ for x' , and we often abuse notation and identify the morphism J with its image in X . By a *partition* of J we will mean a sequence (possibly empty) of integers a_1, \dots, a_n , with $0 < a_1 \leq \dots \leq a_n < d(x, x')$. Setting $y_i = a_i \cdot J$, we will then write

$$x \leq y_1 \leq \dots \leq y_n \leq x'. \quad (3.1)$$

(Although we allow $y_i = y_{i+1}$, later on we will impose conditions which will force $y_{i-1} \neq y_{i+1}$ for $1 < i < n$, and which will force $x \neq y_1$ and $y_n \neq x'$.)

Let a partition of J be given as in (3.1), and let (x_0, \dots, x_{n+1}) be a sequence of points of X , satisfying the following conditions.

- (a) $x_0 = x$ and $x_{n+1} = x'$.
- (b) For all i , $1 \leq i \leq n$, we have:
 - (i) $x_i \notin J$ and $[x_i, y_i]$ is the bridge from x_i to J , and
 - (ii) $y_i = Y(x_{i-1}, x_i, x_{i+1})$.

Define $\Sigma = \Sigma(x_0, \dots, x_{n+1})$ to be the smallest subtree of X containing $\{x_0, \dots, x_{n+1}\}$. Thus:

$$\Sigma = \bigcup_{i,j} [x_i, x_j] = J \cup \left(\bigcup_{i=1}^n [x_i, y_i] \right).$$




FIGURE 3

We say that Σ is a *simplex over J* (relative to the partition $(x \leq y_1 \leq \dots \leq y_n \leq x')$); see Fig. 3.

Recall that \mathcal{H}^{**} denotes the set of all reduced \mathcal{H} -sequences.

3.3. DEFINITION. Let $\mathbf{h} = (h_0, \dots, h_n) \in \mathcal{H}^{**}$, and let $\Sigma = \Sigma(x_0, \dots, x_{n+1})$ be a simplex over J . We say that \mathbf{h} and Σ are *engaged* if, for all i with $0 \leq i \leq n$, we have $\Sigma \cap A(h_i) = [x_i, x_{i+1}]$, oriented toward x_i by h_i .

If \mathbf{h} and Σ are engaged, as shown in Fig. 4, we put

$$J_i = J \cap A(h_i),$$

$$d_i = l(J_i),$$

$$m_i = m(h_i) \quad (\text{see (1.9)}).$$

For any segment $\mathbf{s} = (h_k, \dots, h_l)$ of \mathbf{h} , we put also

$$d(\mathbf{s}) = (d_k, \dots, d_l),$$

$$m(\mathbf{s}) = (m_k, \dots, m_l).$$

3.5. LEMMA. Let $\mathbf{h} = (h_0, \dots, h_n) \in \mathcal{H}^{**}$, and let $\Sigma = \Sigma(x_0, \dots, x_{n+1})$ be a simplex over J . Suppose that \mathbf{h} and Σ are engaged. Then the following hold for all i , $0 < i \leq n+1$.




FIGURE 4

$$(a) \quad \Sigma \cap A(h_{i-1}) \cap A(h_i) = [x_i, y_i].$$

$$(b) \quad d(x_i, y_i) \leq 2, \text{ with strict inequality unless } \{x_i, y_i\} \subseteq D.$$

Proof. Part (a) is by construction. Since $h_{i-1}h_i \notin \mathcal{H} \cup \{1\}$, part (b) then follows from (1.8).

We will be concerned with segments \mathbf{s} of \mathbf{h} which, to begin with, satisfy the following condition.

(*) We have $\mathbf{s} = (h_k, \dots, h_l)$, $0 < k, l < n$, and $l(\Sigma \cap A(h_{i-1}) \cap A(h_i)) = d(x_i, y_i) = 2$ for all $i, k \leq i \leq l + 1$.

3.6. DEFINITION. Let $\mathbf{h} \in \mathcal{H}^{**}$, let $\mathbf{s} = (h_k, \dots, h_l)$ be a segment of \mathbf{h} , and let Σ be a simplex over J . Assume that \mathbf{h} and Σ are engaged.

(I) We say that \mathbf{s} is *special, of type I*, provided that (*) holds, and provided that

$$l(\mathbf{s}) = 1 \text{ or } 3,$$

$$m(\mathbf{s}) = (2) \text{ or } (2, 2, 2),$$

and

$$d(\mathbf{s}) = (4) \text{ or } (2, 4, 2).$$

(II) We say that \mathbf{s} is *special, of type II*, provided that (*) holds, and $m(\mathbf{s})$ is an alternating string of 2's and 3's, and one of the following holds.

$$(i) \quad l(\mathbf{s}) = 1, m(\mathbf{s}) = (3), \text{ and } d(\mathbf{s}) = (6).$$

$$(ii) \quad l(\mathbf{s}) = 2, m(\mathbf{s}) = (2, 3) \text{ or } (3, 2), \text{ and } d(\mathbf{s}) = (4, 4).$$

$$(iii) \quad l(\mathbf{s}) = 3, m(\mathbf{s}) = (2, 3, 2), \text{ and } d(\mathbf{s}) = (4, 2, 4).$$

$$(iv) \quad l(\mathbf{s}) > 4, \text{ and } d(\mathbf{s}) = \lambda \circ \xi_t \circ \mu, \text{ where } t = [(l - k + 1)/4],$$

$$\lambda = \begin{cases} (4, 2) & \text{if } m_k = 3 \\ (4, 2, 2) & \text{if } m_k = 2 \end{cases} \quad \mu = \begin{cases} (2, 4) & \text{if } m_l = 3 \\ (2, 2, 4) & \text{if } m_l = 2, \end{cases}$$

and where ξ_t is defined inductively by

$$\xi_1 = (6), \quad \text{and} \quad \xi_i = \xi_{i-1} \circ (2, 2, 2, 6).$$

3.7. Remark. It follows from (3.6) that $l(\mathbf{s})$ is not divisible by 4. Moreover, if \mathbf{s} is of type II and $l(\mathbf{s}) > 4$, then either

$$l(\mathbf{s}) \equiv 1 \pmod{4}, \quad \text{and} \quad m(\mathbf{s}) = (3, 2, \dots, 2, 3),$$

$$l(\mathbf{s}) \equiv 2 \pmod{4},$$

or

$$l(\mathbf{s}) \equiv 3 \pmod{4}, \quad \text{and} \quad m(\mathbf{s}) = (2, 3, \dots, 3, 2).$$

Next, let \mathbf{x} be a segment of \mathbf{h} of the form

$$\mathbf{x} = (h_{k_1}) \circ \mathbf{s}_1 \circ (h_{k_2}) \circ \cdots \circ \mathbf{s}_u \circ (h_{k_{u+1}}),$$

where each \mathbf{s}_i is special (of type I or II) relative to Σ . Write

$$\mathbf{s}_i = (h_{r_i}, \dots, h_{s_i}), \quad 1 \leq i \leq u,$$

so that $r_i = k_i + 1$ and $s_i = k_{i+1} - 1$. Define integers $p(\mathbf{s}_i)$ and $q(\mathbf{s}_i)$, as

$$\begin{aligned} (p(\mathbf{s}_i), q(\mathbf{s}_i)) &= (2, 2) && \text{if } \mathbf{s}_i \text{ is of type I} \\ (p(\mathbf{s}_i), q(\mathbf{s}_i)) &= (8 - 2 \cdot m_{k_{i+1}}, 8 - 2 \cdot m_{k_{i+1}-1}) && \text{if } \mathbf{s}_i \text{ is of type II.} \end{aligned} \tag{3.8}$$

3.9. DEFINITION. Let \mathbf{h} and Σ be engaged. A segment $\mathbf{x} = (h_{k_1}) \circ \mathbf{s}_1 \circ (h_{k_2}) \circ \cdots \circ \mathbf{s}_u \circ (h_{k_{u+1}})$ of \mathbf{h} will be said to be *extraspecial* (relative to Σ) if $u \geq 1$ and:

- (a) Each \mathbf{s}_i is a special segment of type I or II;
- (b) For all i with $1 \leq i \leq u$, we have $m_{k_i} \geq 6$ and $m_{k_{i+1}} \geq 6$, unless \mathbf{s}_i is of type I and of length 1, in which case we have

$$1/m_{k_i} + 1/m_{k_{i+1}} \leq 1/2;$$

- (c) We have the inequalities

$$l(\Sigma \cap A(h_{k_1})) \geq 2 \cdot m_{k_1} - p(\mathbf{s}_1),$$

$$l(\Sigma \cap A(h_{k_i})) \geq 2 \cdot m_{k_i} - p(\mathbf{s}_1) - q(\mathbf{s}_{i-1}) \quad (1 < i < u + 1),$$

$$l(\Sigma \cap A(h_{k_{u+1}})) \geq 2 \cdot m_{k_{u+1}} - q(\mathbf{s}_u).$$

Moreover, for each i , the corresponding inequality is strict if $\partial(\Sigma \cap A(h_{k_i})) \subseteq D$.

3.10. DEFINITION. A segment \mathbf{a} of \mathbf{h} will be said to be *ordinary* (relative to Σ) if for every segment (h) of \mathbf{a} of length 1 we have

$$l(\Sigma \cap A(h)) \geq 2 \cdot m(h)$$

with strict inequality if $\partial(\Sigma \cap A(h)) \subseteq D$. (In particular, the empty segment (\emptyset) is ordinary.)

3.11. DEFINITION. Let $\mathbf{h} \in \mathcal{H}^{**}$, and let Σ be a simplex over J . We say that \mathbf{h} and Σ are married if \mathbf{h} and Σ are engaged, and if we can write

$$\mathbf{h} = \mathbf{a}_1 \circ \mathbf{x}_1 \circ \cdots \circ \mathbf{a}_m \circ \mathbf{x}_m \circ \mathbf{a}_{m+1},$$

where each \mathbf{a}_i is an ordinary segment, and each \mathbf{x}_j is an estraspecial segment of \mathbf{h} , relative to Σ .

We may now, at long last, define a suitable polarization of X over \mathcal{H} . The reader will need to recall the notion of “distillation” of an \mathcal{H} -sequence, from (2.2).

3.12. DEFINITION. Let J be an oriented closed segment of X , and let $\mathbf{h}' \in \mathcal{H}^*$. We declare

$$J \in \mathcal{P}(\mathbf{h}')$$

if there exists a simplex Σ over J , a braiding \mathbf{h}'' of \mathbf{h}' , and a distillation \mathbf{h} of \mathbf{h}'' , such that \mathbf{h} and Σ are married.

3.13. PROPOSITION. \mathcal{P} is a polarization of X over \mathcal{H} .

Proof. First of all, it is clear from the definitions (3.2) and (3.3) that the only simplices which are engaged with the empty sequence, $(\emptyset) \in \mathcal{H}^*$, are the simplices which consist of a single vertex. Thus, $\mathcal{P}((\emptyset))$ is the set of all degenerate closed segments of X , as required in Definition (2.6). Next, it is immediate from (3.12) that $\mathcal{P}(\mathbf{h}') = \mathcal{P}(\mathbf{h}'')$ for all braidings \mathbf{h}'' of \mathbf{h}' . Fix such a braiding \mathbf{h}'' , and let \mathbf{h} be a distillation of \mathbf{h}'' . Then $\mathbf{h} \in \mathcal{H}^{**}$, by (2.3). For any $g \in \langle \mathcal{H} \rangle$ one sees that \mathbf{h}''^g is a braiding of \mathbf{h}'^g , and \mathbf{h}^g is a distillation of \mathbf{h}''^g . Let $J \in \mathcal{P}(\mathbf{h}')$, and let Σ be a simplex over J such that \mathbf{h} and Σ are married. Then $\Sigma \cdot g$ is a simplex over $J \cdot g$, and it is plain that \mathbf{h}^g and $\Sigma \cdot g$ are married. Thus, $J \cdot g \in \mathcal{P}(\mathbf{h}'^g)$. Finally, (2.4) shows that there is a braiding \mathbf{g} of \mathbf{h}'^{-1} such that \mathbf{h}^{-1} is a distillation of \mathbf{g} . Evidently, \mathbf{h}^{-1} and J^{opp} are married, so we have $J^{opp} \in \mathcal{P}(\mathbf{h}'^{-1})$. Thus, all the requirements of Definition (2.6) have been met by \mathcal{P} .

4. PROPERTIES OF \mathcal{P}

Let \mathcal{P} be the polarization of X over \mathcal{H} which was introduced in the preceding section. Our eventual goal is to show that \mathcal{P} controls convergence relative to (b, \mathcal{D}) , in the sense of Definition (2.8). This will be achieved in Section 5. The aim of the present section is to supply a few supporting results for the arguments in Section 5.

Fix $\mathbf{h}' \in \mathcal{H}^*$, and fix $J \in \mathcal{P}(\mathbf{h}')$. Thus, there is a braiding \mathbf{h}'' of \mathbf{h}' , a distillation \mathbf{h} of \mathbf{h}'' , and a simplex Σ over J , such that \mathbf{h} and Σ are married,

in the sense of (3.11). Write

$$\begin{aligned}
 \mathbf{h} &= (h_0, \dots, h_n), \\
 \Sigma &= \Sigma(x_0, \dots, x_{n+1}), \\
 J &= [x, x'], \quad \text{partitioned by } x \leq y_1 \leq \dots \leq y_n \leq x', \text{ where } x = x_0, \\
 &\quad x' = x_{n+1}, \quad \text{and } y_i = Y(x_{i-1}, x_i, x_{i+1}), \\
 J_i &= J \cap A(h_i), \\
 d_i &= l(J_i), \\
 m_i &= m(h_i).
 \end{aligned}$$

Fix a decomposition of \mathbf{h} as in (3.11),

$$\mathbf{h} = \mathbf{a}_1 \circ \mathbf{x}_1 \circ \mathbf{a}_2 \circ \dots \circ \mathbf{a}_m \circ \mathbf{x}_m \circ \mathbf{a}_{m+1}, \quad (4.1)$$

where each \mathbf{a}_i is an ordinary segment (possibly empty), and each \mathbf{x}_j is an extraspecial segment of \mathbf{h} , relative to Σ . For $\mathbf{x} \in \{\mathbf{x}_1, \dots, \mathbf{x}_m\}$ we can write, as in (3.9),

$$\mathbf{x} = (h_{k_1}) \circ \mathbf{s}_1 \circ (h_{k_2}) \circ \dots \circ \mathbf{s}_u \circ (h_{k_{u+1}}), \quad (4.2)$$

where each \mathbf{s}_i is a special segment (of type I or II) relative to Σ , and where the short segments (h_{k_i}) satisfy the various conditions in (3.9). The segments \mathbf{x}_j , $1 \leq j \leq m$, will be called the *basic extraspecial segments* of \mathbf{h} . A segment \mathbf{s} of \mathbf{h} will be called a *basic special segment* if $\mathbf{s} = \mathbf{s}_i$ in some basic extraspecial segment \mathbf{x} , as in (4.2). We make no attempt to decide whether the decompositions (4.1) and (4.2) are uniquely determined.

We now define a partition $\{\mathcal{S}_j\}_{0 \leq j \leq 3}$ of $\{0, \dots, n+1\}$, as

$$\begin{aligned}
 \mathcal{S}_0 &= \{k : (h_k) \text{ is a segment of some } \mathbf{a}_i\} \\
 \mathcal{S}_1 &= \{k : (h_k) = (h_{k_1}) \text{ or } (h_k) = (h_{k_{u+1}}) \\
 &\quad \text{in some basic extraspecial } \mathbf{x} \text{ as in (4.2)}\} \\
 \mathcal{S}_2 &= \{k : (h_k) = (h_{k_i}), \text{ with } 1 < i < u+1, \\
 &\quad \text{in some basic extraspecial } \mathbf{x} \text{ as in (4.2)}\} \\
 \mathcal{S}_3 &= \{k : (h_k) \text{ is a segment of some basic special segment of } \mathbf{h}\}.
 \end{aligned}$$

We mention that by (h_k) we mean the $(k+1)$ st segment of length 1 in \mathbf{h} . It is entirely possible that $h_j = h_k$ for two distinct indices j and k .

4.3. LEMMA. *Suppose $\mathbf{h}' \neq (\emptyset)$. Then $\mathbf{h} \neq (\emptyset)$ and J is non-degenerate. Indeed, for both $k = 0$ and n , J_k contains at least two points not in D .*

Proof. It is an obvious consequence of Definition (2.2) that $\mathbf{h} \neq (\emptyset)$. Suppose now that J_0 contains at most one point which is not in D . Then $l(J_0) \leq 2$, with equality only if $\partial J_0 = \{x, y_1\} \subseteq D$. Then also (3.5) yields $l(\Sigma \cap A(h_0)) = d(x, x_1) \leq 4$, with equality only if $\{x, x_1\} \subseteq D$. On the other hand, \mathbf{h} and Σ are married, so $0 \in \mathcal{S}_0 \cup \mathcal{S}_1$. If $0 \in \mathcal{S}_0$ then, by definition, we have $d(x, x_1) \geq 2 \cdot m_0$, with strict inequality if $\{x, x_1\} \subseteq D$. Thus, $0 \in \mathcal{S}_1$, and (3.9)(c) gives $d(x, x_1) \geq 2 \cdot m_0 - p(\mathbf{s}_1)$, where \mathbf{s}_1 is the basic special sequence to the right of (h_0) in \mathbf{h} . Again, the inequality is strict if $\{x, x_1\} \subseteq D$. Here (3.9)(b) yields $2 \cdot m_0 - p(\mathbf{s}_1) \geq 6$ unless $m_0 = 3$ and $p(\mathbf{s}_1) = 2$. Thus, $m_0 = 3$, $p(\mathbf{s}_1) = 2$, and $d(x, x_1) = 4$. But then $\{x, x_1\} \subseteq D$ and $d(x, x_1) > 4$, a contradiction.

4.4. LEMMA. *Let $0 \leq k \leq n$ and suppose $d_k \leq 2$. Then $0 < k < n$ and one of the following holds.*

- (i) $k \in \mathcal{S}_0$, $d_k = 2$, and $m_k = 2$.
- (ii) $k \in \mathcal{S}_1$, $d_k = 2$, $m_k = 3$, and the basic special segment to one side of (h_k) is of type I and of length 1.
- (iii) $k \in \mathcal{S}_2$, $d_k = 2$, $m_k = 6$, $m_{k-1} = m_{k+1} = 2$, $d_{k-1} = d_{k+1} = 4$, and the special segments on each side of (h_k) in \mathbf{h} are of type II.
- (iv) $k \in \mathcal{S}_2$, $d_k = 0$ or 2 , and the special segments on each side of (h_k) in \mathbf{h} are of type I and of length 1. If $d_k = 2$ then $m_k = 3$ or 4 , and if $d_k = 0$ then $m_k = 3$.
- (v) $k \in \mathcal{S}_3$, $d_k = 2$, and $m_k = 2$ or 3 . Further, if $m_k = 3$ then $\{k-1, k+1\} \subseteq \mathcal{S}_3$ and $m_{k-1} = m_{k+1} = 2$.

Proof. Suppose first that $k \in \mathcal{S}_0$. Then (3.10) says that $l(\Sigma \cap A(h_k)) = d(x_k, x_{k+1}) \geq 2 \cdot m_k$, with strict inequality if $\{x_k, x_{k+1}\} \subseteq D$. Since $d_k \leq 2$, (3.5) yields $d(x_k, x_{k+1}) \leq 6$, with equality only if $\{x_k, y_k, y_{k+1}, x_{k+1}\} \subseteq D$. Thus, $m_k = 2$ and $d(x_k, x_{k+1}) \geq 4$. If $k = 0$ or n , (3.5) yields $d(x_k, x_{k+1}) = 4$ and $\{x_k, x_{k+1}\} \subseteq D$; a contradiction. Thus, $0 < k < n$. Also, if $d_k < 2$ then (3.5) yields $d(x_k, x_{k+1}) \leq 4$, with equality only if $\{x_k, x_{k+1}\} \subseteq D$. Again, this is inconsistent with (3.10), so $d_k = 2$ and (i) holds.

Suppose next that $k \in \mathcal{S}_1$, so that (h_k) comes either at the beginning or at the end of a basic extraspecial segment \mathbf{x} of \mathbf{h} . By symmetry, we may assume without loss that (h_k) is at the beginning of \mathbf{x} . In the notation of (4.2), $(h_k) = (h_{k_1})$, and \mathbf{s}_1 is the basic special sequence following (h_k) . We apply (3.9)(c) to \mathbf{x} , and obtain $d(x_k, x_{k+1}) \geq 2 \cdot m_k - p(\mathbf{s}_1)$, with strict inequality if $\{x_k, x_{k+1}\} \subseteq D$. Here $p(\mathbf{s}_1)$ is given by (3.8). Thus, $p(\mathbf{s}_1) = 2$ or else \mathbf{s}_1 is of type II and $p(\mathbf{s}_1) = 4$. Since (3.5) gives $d(x_k, x_{k+1}) \leq 6$, we then have $m_k \leq 5$. Then (3.9)(b) says that \mathbf{s}_1 is of type I and length 1, and

$m_k > 2$. Here $p(\mathbf{s}_1) = 2$, so $m_k = 3$ or 4 . Further, if $m_k = 4$ then $d(\mathbf{x}_k, x_{k+1}) = 6$, and (3.5) yields $\{x_k, x_{k+1}\} \subseteq D$, and then (3.9)(c) gives $d(x_k, x_{k+1}) > 6$, a contradiction. Thus, $m_k = 3$, $2 \cdot m_k - p(\mathbf{s}_1) = 4$, and so we have $d(x_k, x_{k+1}) \geq 4$, with strict inequality if $\{x_k, x_{k+1}\} \subseteq D$. This forces $k \neq 0$ (so that the first assertion in (4.4) is now proven), and it forces also $d_k = 2$, in view of (3.5). Thus, (ii) holds.

Suppose next that $k \in \mathcal{S}_2$. Again let \mathbf{x} be the basic extraspecial segment involving (h_k) . In the notation of (4.2) we now have $k = k_i$ for some i , $1 < i < u + 1$, and \mathbf{s}_{i-1} and \mathbf{s}_i are the basic special sequences which, respectively, precede and follow (h_k) in \mathbf{x} . Now (3.9)(c) says

$$d(x_k, x_{k+1}) \geq 2 \cdot m_k - p(\mathbf{s}_i) - q(\mathbf{s}_{i-1}) \quad (*)$$

with the familiar condition on strict inequality. Since also $d(x_k, x_{k+1}) \leq 6$, with strict inequality if $\{x_k, x_{k+1}\} \subseteq D$, we get

$$6 > 2 \cdot m_k - p(\mathbf{s}_i) - q(\mathbf{s}_{i-1}). \quad (**)$$

Since $p(\mathbf{s}_i) \leq 4$ and $q(\mathbf{s}_{i-1}) \leq 4$, by (3.8), we conclude that $m_k \leq 6$.

Suppose $m_k = 6$. Then $p(\mathbf{s}_i) = q(\mathbf{s}_{i-1}) = 4$, and (3.8) says that $m_{k-1} = m_{k+1} = 2$, and both \mathbf{s}_{i-1} and \mathbf{s}_i are of type II, and $d_{k-1} = d_{k+1} = 4$. Now $2 \cdot m_k - p(\mathbf{s}_i) - q(\mathbf{s}_{i-1}) = 4$. If $d_k < 2$ then (3.5) gives $d(x_k, x_{k+1}) \leq 4$, with equality only if $\{x_k, x_{k+1}\} \subseteq D$. Thus, $d_k = 2$ and (iii) holds.

Now suppose $m_k < 6$. Then (3.9)(b) says that \mathbf{s}_{i-1} and \mathbf{s}_i are special of type I and length 1, and (3.8) gives $p(\mathbf{s}_i) = q(\mathbf{s}_{i-1}) = 2$. Also, (3.9)(b) says that $m_k > 2$, and then (**) yields $m_k = 3$ or 4 . If $d_k = 2$ then (iv) holds, so assume $d_k = 0$ or 1 . As we have seen, (3.5) then gives $d(x_k, x_{k+1}) \leq 4$, with equality only if $\{x_k, x_{k+1}\} \subseteq D$. Then (*) yields $m_k = 3$. In order to obtain (iv) it now suffices to show that $d_k \neq 1$. But if $d_k = 1$ then either \mathbf{s}_{i-1} or \mathbf{s}_i fails to satisfy the condition (*) preceding, and involved in, Definition (3.6). Thus, $d_k \neq 1$ and (iv) holds.

Finally, assume that $k \in \mathcal{S}_3$. Then (3.6) says that $d_k = 2$ and $m_k = 2$ or 3 . Assume further that $m_k = 3$, and let \mathbf{s} be the basic special segment in \mathbf{h} involving (h_k) . Then \mathbf{s} is of type II, and inspection of (3.6) reveals that, with $d_k = 2$, h_k does not occur as either the first or last term in the sequence \mathbf{s} . Thus, we have $k \pm 1 \in \mathcal{S}_3$ and $m_{k \pm 1} = 2$, so that (v) holds.

4.5. LEMMA. *Assume that $d_k = d_{k+1} = 2$, and that $m_k + m_{k+1} > 4$. Then one of the following holds.*

(i) $k \in \mathcal{S}_1$, $k + 1 \in \mathcal{S}_0$, $m_k = 3$, $m_{k+1} = 2$, and (h_{k-1}) is a basic special segment of type I.

(ii) $k \in \mathcal{S}_0$, $k + 1 \in \mathcal{S}_1$, $m_k = 2$, $m_{k+1} = 3$, and (h_{k+2}) is a basic special segment of type I.

(iii) $\{k, k+1\} \subseteq \mathcal{S}_1$, $m_k = m_{k+1} = 3$, and both (h_{k-1}) and (h_{k+2}) are basic special segments of type I.

(iv) $\{k, k+1\} \subseteq \mathcal{S}_3$, and (h_k, h_{k+1}) is part of a basic special segment of type II.

Proof. Suppose $k \in \mathcal{S}_0$. If also $k+1 \in \mathcal{S}_0$ then $m_k + m_{k+1} = 4$ by (4.4)(i). Thus, we must have $k+1 \in \mathcal{S}_1$, and then (ii) follows from outcomes (i) and (ii) of (4.4). Similarly, if $k+1 \in \mathcal{S}_0$ then we get (i). So, we may now assume $k \notin \mathcal{S}_0$ and $k+1 \notin \mathcal{S}_0$. If $\{k, k+1\} \subseteq \mathcal{S}_1$ then (iii) follows from (4.4)(ii). If $k \in \mathcal{S}_2$ then $d_{k+1} = 4$ by (4.4)(iii), so we now conclude that $\{k, k+1\} \subseteq \mathcal{S}_3$. Then (h_k, h_{k+1}) is part of a basic special sequence \mathbf{s} in \mathbf{h} . Since $m_k + m_{k+1} > 4$, \mathbf{s} is then of type II, and (iv) holds.

4.6. LEMMA. Assume that $d_{k-1} = d_k = d_{k+1} = 2$, and assume that both $m_{k-1} + m_k > 4$ and $m_k + m_{k+1} > 4$. Then one of the following holds.

(i) $k \in \mathcal{S}_0$, $\{k-1, k+1\} \subseteq \mathcal{S}_1$, $m_{k-1} = m_{k+1} = 3$, and both (h_{k-2}) and (h_{k+2}) are basic special segments of type I.

(ii) $\{k-2, k-1, k, k+1, k+2\} \subseteq \mathcal{S}_3$, $d_{k-2} = d_{k+2} = 6$, $m_{k-1} = m_{k+1} = 2$, and $m_{k-2} = m_k = m_{k+2} = 3$.

Proof. If $k \in \mathcal{S}_0$ we get (i) as a consequence of (4.5)(i), (ii). If $k \in \mathcal{S}_1$ then (4.5) shows that $\{k-1, k+1\} \subseteq \mathcal{S}_0 \cup \mathcal{S}_1$, which is contrary to the definition of \mathcal{S}_1 . Now (4.5) forces $k \in \mathcal{S}_3$, and then also (h_{k-1}, h_k, h_{k+1}) is part of a basic special sequence \mathbf{s} of type II. Since $d_{k-1} = d_k = d_{k+1} = 2$, a glance at (3.6)(II) shows that $d_{k-2} = d_{k+2} = 6$, $m_{k-2} = m_{k+2} = 3$, and $(h_{k-2}, \dots, h_{k+2})$ is part of \mathbf{s} . Thus, (ii) holds.

For the remainder of this section, and in the following section, we fix an element h of $\mathcal{E}(\mathbf{h}', J)$. Recall from Definition (2.7) that this means that $\mathbf{h}' \circ (h) \in \mathcal{H}^*$, and $J \cap A(h)$ is a closed segment, oriented toward x' by h , and non-degenerate if J is non-degenerate. Put

$$I = J \cap A(h) = [u, v]$$

with $[u, v] \leq [x, x']$. (That is to say, $d(u, x) \leq d(v, x)$ and $d(u, x') \geq d(v, x')$.) Thus, I is oriented toward v by h . Put

$$\mathcal{S}_4 = \{k : y_k \in [u, v], u \neq y_k \neq v, 1 \leq k \leq n\}.$$

That is, $k \in \mathcal{S}_4$ if y_k is in the “interior” of I .

4.7. LEMMA. For any k with $0 \leq k \leq n$, $A(h) \cap A(h_k)$ contains at most one vertex not in D . In particular, $l(I \cap J_k) \leq 2$.

Proof. This is immediate from (2.3) and (1.8).

4.8. LEMMA. *Let $k \in \mathcal{S}_4$, and suppose $y_k \neq y_{k+1}$. Then $1/m_k + 1/m_{k+1} + 1/m(h) \leq 1$.*

Proof. This is immediate from (2.3) and (1.10).

4.9. LEMMA. *Suppose $l(I) \geq \frac{1}{2}a(h)$. Then $\partial J \cap I = \emptyset$, $\{u, y_r, \dots, y_s, v\} \subseteq D$, and one of the following holds.*

(i) $\mathcal{S}_4 = \{r\}$, $a(h) = 8$, and $d(u, y_r) = d(y_r, v) = 2$.

(ii) $\mathcal{S}_4 = \{r, r+1\}$, $y_r = y_{r+1}$, $a(h) = 8$, and $d(u, y_r) = d(y_{r+1}, v) =$

2.

(iii) $\mathcal{S}_4 = \{r, r+1\}$, $y_r \neq y_{r+1}$, $a(h) = 12$, and $d(u, y_r) = d(y_r, y_{r+1}) = d(y_{r+1}, v) = 2$.

Moreover, we either have $r \in \mathcal{S}_0$ and $m_r = 2$, or else $r \in \mathcal{S}_2$ and $m_r = 6$.

Proof. Suppose $x \in I$. As $l(I) > 2$, it then follows from (4.3) that $I \cap J_0$ contains two points not in D , and this contradicts (4.7). Thus, $x \notin I$, and similarly $x' \notin I$. The same reasoning shows also that $I \not\subseteq J_k$ for any k , $0 \leq k \leq n$, and hence $\mathcal{S}_4 \neq \emptyset$.

Suppose next that $y_k = y_{k+1}$ for some k , with $r \leq k < s$. Then $d_k = 0$, so that (h_{k-1}) and (h_{k+1}) are basic special sequences, of type I, by (4.4). Consulting Definition (3.6), this means that $d_{k-1} = d_{k+1} = 4$. Then (4.7) forces $u \in J_{k-1}$, $v \in J_{k+1}$, with each of $[u, y_k]$ and $[y_{k+1}, v]$ containing at most one point not in D . Since $a(h) \geq 8$, by (1.12), it then follows that $a(h) = 8$, $m(h) = 2$, $d(u, y_k) = d(y_{k+1}, v) = 2$, and $\{u, y_k, v\} \subseteq D$. Thus, (ii) holds in this case.

Suppose next that $r = s$. Thus, $u \in J_{r-1}$ and $v \in J_r$. As in the preceding paragraph, each of $[u, y_k]$ and $[y_k, v]$ contains at most one point not in D , and since $a(h) \geq 8$, (i) holds in this case.

We may now assume $r < s$, and $y_k \neq y_{k+1}$ for any k with $r \leq k < s$. Then (4.7) and (4.4) together yield $d_r = \dots = d_{s-1} = 2$, and $\{y_r, \dots, y_s\} \subseteq D$. Then also (4.8) yields

$$1/m_k + 1/m_{k+1} + 1/m(h) \leq 1 \quad (**)$$

for all k with $r \leq k < s$. In particular, we have $m_k + m_{k+1} > 4$ for all such k .

Suppose $s - r \geq 3$, and take $k = r + 1$, so that $\{k - 1, \dots, k + 2\} \subseteq \mathcal{S}_4$, and we are in one of the two situations described in (4.6). Suppose (4.6)(i) applies. Thus, $k \in \mathcal{S}_0$, $\{k \pm 1\} \subseteq \mathcal{S}_1$, (h_{k-2}) and (h_{k+2}) are basic special segments of type I, and $m_{k-1} = m_{k+1} = 3$. Consulting (3.6) we see that $d_{k-2} = d_{k+2} = 4$, and $m_{k-2} = m_{k+2} = 2$. Apply (**) with $k - 2$ in place of k , and conclude that $m(h) \geq 6$. Then $\frac{1}{2}a(h) \geq 12$, and this implies $l(I \cap J_{k-2}) \geq 3$ or $l(I \cap J_{k+2}) \geq 3$, which is contrary to (4.7).

Now (4.6)(ii) must hold, so that $d_{k-2} = d_{k+2} = 6$, $m_{k-1} = m_{k+1} = 2$, and $m_{k-2} = m_k = m_{k+2} = 3$. Again, $(**)$ yields $m(h) \geq 6$ and $l(I) \geq 12$, with a contradiction as in the preceding paragraph. We therefore conclude that $s - r \leq 2$.

Suppose $s - r = 2$. Then $d(y_r, y_s) \leq 4$ and $l(I) \leq 8$, so that $m(h) \leq 4$. Then $(**)$ yields $m_r + m_{r+1} > 5$. We may appeal to (4.5), with $k = r$, where we see that the condition $m_r + m_{r+1} > 5$ immediately rules out all but (4.5)(iii). But in that case we have $m_{r-1} = 2$, and applying $(**)$, with $r - 1$ in place of k , again yields a contradiction. We conclude that $s = r + 1$.

Now $d(y_r, y_s) = 2$ and $l(I) \geq 6$, so that $a(h) = 8$ or 12 . Suppose $a(h) = 8$. Then $(**)$ yields $m_k > 2$ for $k \in \{r - 1, r, r + 1\}$. But $d_r = 2$, so (4.4) provides a contradiction. Thus, $a(h) = 12$ and hence $l(I) = 6$. Further, (4.7) yields $\{u, y_r, y_{r+1}, v\} \subseteq D$ and $d(u, y_r) = d(y_r, y_{r+1}) = d(y_{r+1}, v) = 2$. Since $(**)$ holds for both $k = r$ and $k = r + 1$, we may exclude the cases given by parts (ii), (iv), and (v) of (4.4), with $k = r$. Thus (4.4)(i) or (iii) applies, and this yields outcome (iii) of the lemma, as desired.

5. CONTROL OF CONVERGENCE

We now show that the polarization \mathcal{P} , defined in (3.12), controls convergence relative to (b, \mathcal{D}) .

We continue with all of the notation introduced in the preceding section. In particular, we have $J \in \mathcal{P}(\mathbf{h}')$, \mathbf{h} a distillation of a braiding \mathbf{h}'' of \mathbf{h}' , and Σ a simplex over J such that \mathbf{h} and Σ are married. Further, we have $h \in \mathcal{E}(\mathbf{h}', J)$.

5.1. LEMMA. *\mathcal{P} satisfies the “Exclusion” condition (2.8)(1).*

Proof. Suppose $\mathbf{h}' \neq \emptyset$. Then (4.3) and (4.7) together imply that $J_0 \not\subseteq A(h)$. Recall the notation: $I = J \cap A(h)$, and notice that an immediate consequence of (4.9) is that $l(I) \leq \frac{1}{2}a(h)$. Finally, suppose $\partial J \cap I \neq \emptyset$. Then (4.3) and (4.7) yield $l(I) \leq 2$. Since $b(h) \geq 8$, by (1.12), we then have $b(h) > 2 \cdot l(I)$, and all pats of (2.8)(1) have been verified.

The problem now is to verify that \mathcal{P} satisfies the “Extension” condition (2.8)(2). To do this, we first handle the cases where $\partial J \cap I \neq \emptyset$. In particular, we need to examine the case where J is degenerate.

5.2. LEMMA. *Let K be a segment of $A(h)$. Assume that $l(K) \geq \frac{1}{2}b(h)$, with strict inequality if $\partial K \subseteq D$. Then $K \in \mathcal{P}((h))$.*

Proof. Notice that (h) is the unique distillation of the unique braiding of (h) , so it suffices to show that (h) and K are married. In terms of the

decomposition (4.1) we take $(h) = \mathbf{a}_1$ and we obtain $l(K \cap A(h)) \geq 2 \cdot m(h)$ (with strict inequality if $\partial K \subseteq D$), since $m(h) = \frac{1}{2}b(h)$. Thus (h) is an ordinary segment of (h) , relative to K , in the sense of definition (3.10). Then (h) and K are married, and thus $K \in \mathcal{P}((h))$.

5.3. LEMMA. *Suppose that we have a point $w \in A(h)$, with $x \in [w, w \cdot h]$. Suppose further that $d(x, w \cdot h) \geq 2 \cdot m(h)$, and that strict inequality holds if $\{x, w\} \subseteq D$. Put $\tilde{J} = [w \cdot h, x']$. Then $\tilde{J} \in \mathcal{P}(\mathbf{h}' \circ (h))$.*

Proof. By assumption we have $x \in I$. If $J = \{x\}$ then $\mathbf{h} = (\emptyset)$ by (4.3), and then $\tilde{J} \in \mathcal{P}((h))$ by (5.2). Thus, we may assume that J is non-degenerate, whence also $\mathbf{h} \neq (\emptyset)$. Now (4.3) and (4.7) together show that $I = [x, v] \leq J_0$. Moreover, (4.7) yields $d(x, v) \leq 2$, with strict inequality unless $\{x, v\} \subseteq D$.

Put $\tilde{\mathbf{h}} = (h) \circ \mathbf{h}$, and notice that $\tilde{\mathbf{h}}$ is a simple distillation of $\mathbf{h} \circ (h)$, of codegree 1, as defined in (2.1). Hence $\tilde{\mathbf{h}}$ is a distillation of $\mathbf{h}' \circ (h)$, which is in turn a braiding of $\mathbf{h}' \circ (h)$. Next, define points $\tilde{x}_0, \dots, \tilde{x}_{n+2}$ by

$$\tilde{x}_0 = w \cdot h, \quad \tilde{x}_i = x_{i-1} \quad (1 \leq i \leq n+2).$$

Put $\tilde{\Sigma} = \Sigma(\tilde{x}_0, \dots, \tilde{x}_{n+2})$. Also, put

$$\tilde{y}_1 = v, \quad \tilde{y}_i = y_{i-1} \quad (2 \leq i \leq n+1).$$

Then $\tilde{\Sigma}$ is a simplex over \tilde{J} , relative to the partition $(\tilde{x}_0 < \tilde{y}_1 \leq \dots \leq \tilde{y}_{n+1} < \tilde{x}_{n+2})$. To see this, we need only sketch a picture, as shown in Fig. 5. Moreover, Fig. 5 displays that $\tilde{\mathbf{h}}$ and $\tilde{\Sigma}$ are engaged. In order to complete the proof of (5.3), it now suffices to show that $\tilde{\mathbf{h}}$ and $\tilde{\Sigma}$ are married.

Write $\mathbf{h} = \mathbf{a}_1 \circ \mathbf{x}_1 \circ \dots \circ \mathbf{a}_m \circ \mathbf{x}_m \circ \mathbf{a}_{m+1}$ as in (4.1). Put $\tilde{\mathbf{a}}_1 = (h) \circ \mathbf{a}_1$, $\tilde{\mathbf{x}}_i = \mathbf{x}_i$ for all i , and $\tilde{\mathbf{a}}_j = \mathbf{a}_j$ for $j > 1$. It is immediate that each $\tilde{\mathbf{x}}_i$ is




FIGURE 5

extraspecial and that each \tilde{a}_j ($j > 1$) is ordinary, relative to $\tilde{\Sigma}$. But also, we have $d(\tilde{x}_0, \tilde{x}_1) = d(w \cdot h, x) \geq 2 \cdot m(h)$, with strict inequality if $\{x, w\} \subseteq D$. Here $w \in D$ if and only if $w \cdot h \in D$, since $a(h)$ is even. Thus $l(\tilde{\Sigma} \cap A(h)) \geq 2 \cdot m(h)$, with strict inequality if $(\tilde{\Sigma} \cap A(h)) \subseteq D$. This shows that $\tilde{\mathbf{a}}_1$ is ordinary, and so $\tilde{\mathbf{h}}$ and $\tilde{\Sigma}$ are married.

Remark. With (5.3), we have shown that part (b) of the Extension condition (2.8)(2) is satisfied by \mathcal{P} .

In order to complete the proof that \mathcal{P} controls convergence relative to (b, \mathcal{D}) , it only remains to deal with part (a) of the Extension condition (2.8)(2). Thus, we may now assume:

5.4. We have $\partial J \cap A(h) = \emptyset$, and $l(I) \geq a(h) - 2 \cdot m(h)$, with strict inequality if $\partial I \not\subseteq D$.

Our goal now will be to show that $[x \cdot h, x'] \in \mathcal{P}(\mathbf{h}' \circ (h))$.

Recall the notation from Section 4: we have $I = [u, v] \leq [x, x']$, and we have $\mathcal{S}_4 = \{r, r+1, \dots, s\}$, defined by $k \in \mathcal{S}_4$ if $y_k \in I - \{u, v\}$. Put

$$\tilde{\mathbf{h}} = (h_0, \dots, h_{r-1})^h \circ (h) \circ (h_s, \dots, h_n),$$

and notice that $\tilde{\mathbf{h}}$ is a simple distillation of $\mathbf{h} \circ (h)$, of codegree 1. Then $\tilde{\mathbf{h}}$ is a distillation of $\mathbf{h}' \circ (h)$, which is itself a braiding of $\mathbf{h}' \circ (h)$. (See (2.1) and (2.2).) Put

$$\tilde{J} = [x \cdot h, x'].$$

We will show that there exists a suitable simplex $\tilde{\Sigma}$ over \tilde{J} , such that $\tilde{\mathbf{h}}$ and $\tilde{\Sigma}$ are married.

There are three cases to consider, corresponding to the three possible outcomes in (4.9). We will treat each case separately.

5.5. LEMMA. Suppose $\mathcal{S}_4 = \{r\}$, $a(h) = 8$, and $d(u, y_r) = d(y_r, v) = 2$. Then $\tilde{J} \in \mathcal{P}(\mathbf{h}' \circ (h))$.

Proof. Define points \tilde{x}_i , $0 \leq i \leq n+2$, as

$$\tilde{x}_0 = x_0 \cdot h, \dots, \tilde{x}_{r-1} = x_{r-1} \cdot h,$$

$$\tilde{x}_r = y_r \cdot h, \quad \tilde{x}_{r+1} = y_r,$$

$$\tilde{x}_{r+2} = x_{r+1}, \dots, \tilde{x}_{n+2} = x_{n+1}.$$

Put $\tilde{\Sigma} = \Sigma(\tilde{x}_0, \dots, \tilde{x}_{n+2})$. Also define points \tilde{y}_i , $1 \leq i \leq n+1$, as

$$\tilde{y}_1 = y_1 \cdot h, \dots, \tilde{y}_{r-1} = y_{r-1} \cdot h,$$

$$\tilde{y}_r = u \cdot h, \quad \tilde{y}_{r+1} = v,$$

$$\tilde{y}_{r+2} = y_{r+1}, \dots, \tilde{y}_{n+1} = y_n.$$

We have the picture of Σ shown in Fig. 6.




FIGURE 6

Since $a(h) = 8$, and since $[u, v]$ is oriented toward v by h (in accordance with (2.7)(c)), we then have the picture of $\tilde{\Sigma}$ shown in Fig. 7.

Evidently, $\tilde{\Sigma}$ is a simplex over \tilde{J} , relative to the partition $(\tilde{x}_0 < \tilde{y}_1 \leq \dots \leq \tilde{y}_{n+1} < \tilde{x}_{n+2})$ of \tilde{J} . Further, it is evident that $\tilde{\mathbf{h}}$ and $\tilde{\Sigma}$ are engaged. It remains to show that $\tilde{\mathbf{h}}$ and $\tilde{\Sigma}$ are married.

Suppose $\{r-1, r\} \subseteq \mathcal{S}_0$. In terms of the decomposition (4.1), this means that (h_{r-1}, h_r) is part of an ordinary segment \mathbf{a}_k of \mathbf{h} , $1 \leq k \leq m+1$. Put $\mathbf{x} = (h_{r-1}^h, h_r)$. We claim that \mathbf{x} is an extraspecial segment of $\tilde{\mathbf{h}}$, relative to $\tilde{\Sigma}$. Indeed, first notice that (h) is a special segment of type I, as is revealed by a glance at the definition (3.6)(I). Then notice that (4.8) yields

$$1/m(h_{r-1}^h) + 1/m(h_r) \leq 1/2,$$

so that \mathbf{x} satisfies (3.9)(b). Since $r-1 \in \mathcal{S}_0$, (3.10) gives $d(x_{r-1}, x_r) \geq 2 \cdot m_r$ with strict inequality if $\{x_{r-1}, x_r\} \subseteq D$. Also $d(y_r, x_r) \leq 2$, with strict inequality unless $\{x_r, y_r\} \subseteq D$, by (3.5). But $l(\tilde{\Sigma} \cap A(h_{r-1}^h)) \geq d(x_{r-1} \cdot h, y_r \cdot h) = d(x_{r-1}, y_r)$, so we get

$$l(\tilde{\Sigma} \cap A(h_{r-1}^h)) \geq 2 \cdot m_{r-1} - 2,$$

with strict inequality if $\partial(\tilde{\Sigma} \cap A(h_{r-1}^h)) \subseteq D$. Since $p((h)) = 2$, according to (3.8), we now see that h_{r-1}^h satisfies (3.9)(c) with respect to \mathbf{x} and $\tilde{\Sigma}$. A symmetric argument yields also (3.9)(c) for h_r . Thus, \mathbf{x} is extraspecial.




FIGURE 7

Now write $\mathbf{a}_k = \mathbf{b}_k \circ (h_{r-1}, h_r) \circ \mathbf{c}_k$. We then have

$$\tilde{\mathbf{h}} = \mathbf{a}_1^h \circ \mathbf{x}_1^h \circ \cdots \circ \mathbf{x}_{k-1}^h \circ \mathbf{b}_k^h \circ \mathbf{x} \circ \mathbf{c}_k \circ \mathbf{x}_k \circ \cdots \circ \mathbf{x}_m \circ \mathbf{a}_{m+1},$$

and this alternating sequence of ordinary and extraspecial segments shows that $\tilde{\mathbf{h}}$ and $\tilde{\Sigma}$ are married in this case.

Suppose next that $r-1 \in \mathcal{S}_0$ and $r \in \mathcal{S}_1$. Thus, (h_{r-1}) comes at the end of an ordinary segment \mathbf{a}_k , and (h_r) comes at the beginning of a basic extraspecial segment \mathbf{x}_k of \mathbf{h} . Put $\mathbf{x} = (h_{r-1}^h, h) \circ \mathbf{x}_k$. We will now show that \mathbf{x} is an extraspecial segment of $\tilde{\mathbf{h}}$, relative to $\tilde{\Sigma}$. Indeed, the proof is almost the same as in the preceding case. Thus, we observe that (h) is a special segment of type I, and $1/m(h_{r-1}^h) + 1/m(h_r) \leq \frac{1}{2}$, so that \mathbf{x} satisfies (3.9)(b). With (h) of type I we have $p((h)) = q((h)) = 2$, by (3.8). Let \mathbf{s} be the basic special segment which follows (h_r) in \mathbf{x}_k . Applying (3.9)(c) to \mathbf{x}_k , we have $d(x_r, x_{r+1}) \geq 2 \cdot m_r - p(\mathbf{s})$, with strict inequality if $\{x_r, x_{r+1}\} \subseteq D$. Now $d(\tilde{x}_{r+1}, \tilde{x}_{r+2}) = d(y_r, x_{r+1}) \geq d(x_r, x_{r+1}) - 2$, with strict inequality unless $\{x_r, y_r\} \subseteq D$, by (3.5). Thus, we get

$$l(\tilde{\Sigma} \cap A(h_r)) \geq 2 \cdot m_r - p(\mathbf{s}) - q((h)),$$

with strict inequality if $\{y_r, x_{r+1}\} \subseteq D$. This is half of what we need in order to show that \mathbf{x} satisfies (3.9)(c). The other half is

$$l(\tilde{\Sigma} \cap A(h_{r-1}^h)) \geq 2 \cdot m_{r-1} - p((h)),$$

which is proved by exactly the same argument as in the case $\{r-1, r\} \subseteq \mathcal{S}_0$, above. Thus, \mathbf{x} is extraspecial relative to $\tilde{\Sigma}$.

Now write $\mathbf{a}_k = \mathbf{b}_k \circ (h_{r-1})$, and obtain

$$\tilde{\mathbf{h}} = \mathbf{a}_1^h \circ \mathbf{x}_1^h \circ \cdots \circ \mathbf{x}_{k-1}^h \circ \mathbf{b}_k^h \circ \mathbf{x} \circ \mathbf{a}_{k+1} \circ \cdots \circ \mathbf{x}_m \circ \mathbf{a}_{m+1}.$$

This alternating sequence of ordinary and extraspecial segments shows that $\tilde{\mathbf{h}}$ and $\tilde{\Sigma}$ are married in this case. Almost the same argument handles the case where $r-1 \in \mathcal{S}_1$ and $r \in \mathcal{S}_0$, so we may assume henceforth that neither $r-1$ nor r is in \mathcal{S}_0 .

Suppose next that $\{r-1, r\} \subseteq \mathcal{S}_1$. Thus, there are basic extraspecial segments \mathbf{x}_{k-1} and \mathbf{x}_k of \mathbf{h} such that \mathbf{x}_{k-1} ends in (h_{r-1}) , \mathbf{x}_k begins in (h_r) , and the ordinary segment \mathbf{a}_k is empty. In this case we put $\mathbf{x} = \mathbf{x}_{k-1}^h \circ (h) \circ \mathbf{x}_k$ and we show that \mathbf{x} is extraspecial relative to $\tilde{\Sigma}$. Let \mathbf{s} be the basic special segment which follows (h_r) in \mathbf{x}_k , and let \mathbf{t} be the basic special segment which precedes (h_{r-1}) in \mathbf{x}_{k-1} . Again, (h) is special of type I, and \mathbf{x} satisfies (3.9)(b), so we need to show

$$l(\tilde{\Sigma} \cap A(h_{r-1}^h)) \geq 2 \cdot m_{r-1} - p((h)) - q(\mathbf{t}),$$

$$l(\tilde{\Sigma} \cap A(h_r)) \geq 2 \cdot m_r - p(\mathbf{s}) - q((h)),$$

and with the familiar requirements concerning strict inequality. But the argument for $\tilde{\Sigma} \cap A(h_r)$ has already been provided in the case $r - 1 \in \mathcal{S}_0$, $r \in \mathcal{S}_1$, considered above. The argument for $\tilde{\Sigma} \cap A(h_{r-1}^h)$ may then safely be deleted, since it is essentially the same as that for $\tilde{\Sigma} \cap A(h_r)$. Thus, \mathbf{x} is extraspecial relative to $\tilde{\Sigma}$, and then $\tilde{\mathbf{h}}$ and $\tilde{\Sigma}$ are married, as is seen from the decomposition

$$\tilde{\mathbf{h}} = \mathbf{a}_1^h \circ \mathbf{x}_1^h \circ \cdots \circ \mathbf{a}_{k-1}^h \circ \mathbf{x} \circ a_{k+1} \circ \cdots \circ \mathbf{x}_m \circ \mathbf{a}_{m+1}.$$

If $\{r - 1, r\} \subseteq \mathcal{S}_3$ then (3.6) shows that $m_{r-1} = 2$ or $m_r = 2$. But $m(h) = 2$, so we contradict (4.8) in this case. By symmetry, we may therefore assume henceforth that $r - 1 \in \mathcal{S}_1 \cup \mathcal{S}_2$ and $r \in \mathcal{S}_3$. Thus, there is a basic extraspecial segment \mathbf{x} of \mathbf{h} which, in the notation of (4.2), satisfies $(h_{r-1}) = (h_{k_i})$ for some i , $1 \leq i \leq u$. Write $\mathbf{x} = \mathbf{y} \circ \mathbf{s}_i \circ \mathbf{z}$, where \mathbf{s}_i is the basic special segment that begins with (h_r) . Then put $\tilde{\mathbf{s}} = (h) \circ \mathbf{s}_i$ and put $\tilde{\mathbf{x}} = \mathbf{y}^h \circ \tilde{\mathbf{s}} \circ \mathbf{z}$. We will now show that $\tilde{\mathbf{s}}$ is a special segment, and $\tilde{\mathbf{x}}$ an extraspecial segment, of $\tilde{\mathbf{h}}$ relative to $\tilde{\Sigma}$.

To see that $\tilde{\mathbf{s}}$ is special, notice first of all that since $d(y_r \cdot h, u \cdot h) = d(y_r, v) = 2$, $\tilde{\mathbf{s}}$ satisfies the condition $(*)$ preceding (3.6). Next, since $r \in \mathcal{S}_3$ we have $m_r = 2$ or 3 , and then (4.8) forces $m_r = 3$ and $m_{r-1} \geq 6$. With $m_r = 3$ we conclude that \mathbf{s}_i is special of type II, and $m(\tilde{\mathbf{s}})$ is an alternating string of 2's and 3's.

Notice that $l(\tilde{J} \cap A(h_r)) = d_r - 2$, since $[y_r, v]$ is now part of $\tilde{\Sigma}$ but not of \tilde{J} . (See the pictures above.) Suppose $l(\mathbf{s}_i) = 1$. Then (3.6)(II) gives $d_r = 6$, so with respect to $\tilde{\Sigma}$ we get $m((h, h_r)) = (2, 3)$ and $d((h, h_r)) = (4, 4)$, as in (3.6)(II)(ii). Thus, $\tilde{\mathbf{s}}$ is special of type II in this case. Similarly, if $l(\mathbf{s}_i) = 2$ then $m(\mathbf{s}_i) = (3, 2)$ and $d(\mathbf{s}_i) = (4, 4)$, and with respect to $\tilde{\Sigma}$ we get $m(\tilde{\mathbf{s}}) = (2, 3, 2)$ and $d(\tilde{\mathbf{s}}) = (4, 2, 4)$. Again, $\tilde{\mathbf{s}}$ is special of type II. According to (3.6)(II), it remains to consider the case where $l(\mathbf{s}_i) > 4$, and where $d(\mathbf{s}_i) = (4, 2) \circ \xi_t \circ \mu$, with $t = [l(\mathbf{s}_i)/4]$. By the remark following (3.6), $l(\mathbf{s}_i) \not\equiv 3 \pmod{4}$, so that $t = [l(\tilde{\mathbf{s}})/4]$. Now $d(\tilde{\mathbf{s}}) = (4, 2, 2) \circ \xi_t \circ \mu$, and thus $\tilde{\mathbf{s}}$ is special of type II.

We next show that $\tilde{\mathbf{x}}$ is extraspecial. We have the decomposition

$$\tilde{\mathbf{x}} = ((h_{k_1}) \circ \mathbf{s}_1 \circ \cdots \circ \mathbf{s}_{i-1} \circ (h_{k_i}))^h \circ \tilde{\mathbf{s}} \circ (h_{k_{i+1}}) \circ \cdots \circ \mathbf{s}_{k_u} \circ (h_{k_{u+1}}).$$

This is a chain of special sequences relative to $\tilde{\Sigma}$, together with “connecting elements” for which we must verify the conditions in (3.9). Notice that (3.9)(b) holds since \mathbf{s}_i is of type II, forcing $m(h_{k_i}) \geq 6$ and $m(h_{k_{i+1}}) \geq 6$. It only remains to show that

$$l(\tilde{\Sigma} \cap A(h_{r-1}^h)) \geq 2 \cdot m_{r-1} - p(\tilde{\mathbf{s}}) - q(\mathbf{s}_{i-1}^h),$$

with strict inequality if $\{x_{r-1} \cdot h, y_r \cdot h\} \subseteq D$, and where we take $q(\mathbf{s}_{i-1}^h) = 0$ if $i = 1$. Here, (3.8) gives $p(\tilde{\mathbf{s}}) = 4$ since $\tilde{\mathbf{s}}$ is of type II, and since $\tilde{\mathbf{s}}$ begins with (h) , where $m(h) = 2$. On the other hand, $p(\mathbf{s}_i) = 2$, also by (3.8). Now $q(\mathbf{s}_{i-1}^h) = q(\mathbf{s}_{i-1})$, so if we apply (3.9)(c) to \mathbf{x} then we get

$$l(\Sigma \cap A(h_{r-1})) \geq 2 \cdot m_{r-1} - p(\tilde{\mathbf{s}}) - q(\mathbf{s}_{i-1}^h) + 2,$$

with strict inequality if $\{x_{r-1}, x_r\} \subseteq D$. But $l(\tilde{\Sigma} \cap A(h_{r-1}^h)) = l(\Sigma \cap A(h_{r-1})) - d(y_r, x_r)$. Since $d(y_r, x_r) \leq 2$, with equality only if $\{x_r, y_r\} \subseteq D$, we have the desired result for $\tilde{\Sigma} \cap A(h_{r-1}^h)$.

Now $\tilde{\mathbf{x}}$ is extraspecial. Taking \mathbf{x} to be \mathbf{x}_k in the decomposition (4.1), we have

$$\tilde{\mathbf{h}} = \mathbf{a}_1^h \circ \mathbf{x}_1^h \circ \cdots \circ \mathbf{a}_k^h \circ \tilde{\mathbf{x}} \circ \mathbf{a}_{k+1} \circ \cdots \circ \mathbf{x}_m \circ \mathbf{a}_{m+1},$$

and evidently $\tilde{\mathbf{h}}$ and $\tilde{\Sigma}$ are married. This completes the proof of (5.5).

5.6. LEMMA. *Suppose $\mathcal{S}_4 = \{r, r+1\}$, $y_r = y_{r+1}$, $a(h) = 8$, and $d(u, y_r) = d(y_{r+1}, v) = 2$. Then $\tilde{J} \in \mathcal{P}(\mathbf{h}' \circ (h))$.*

Proof. Define points \tilde{x}_i , $0 \leq i \leq n+1$, as

$$\begin{aligned} \tilde{x}_0 &= x_0 \cdot h, \dots, \tilde{x}_{r-1} = x_{r-1} \cdot h, \\ \tilde{x}_r &= y_r \cdot h, \quad \tilde{x}_{r+1} = y_r = y_{r+1}, \\ \tilde{x}_{r+2} &= x_{r+2}, \dots, \tilde{x}_{n+1} = x_{n+1}. \end{aligned}$$

Put $\tilde{\Sigma} = \Sigma(\tilde{x}_0, \dots, \tilde{x}_{n+1})$. Also, define points \tilde{y}_j , $1 \leq j \leq n$, as

$$\begin{aligned} \tilde{y}_1 &= y_1 \cdot h, \dots, \tilde{y}_{r-1} = y_{r-1} \cdot h, \\ \tilde{y}_r &= u \cdot h, \quad \tilde{y}_{r+1} = v, \\ \tilde{y}_{r+2} &= y_{r+2}, \dots, \tilde{y}_n = y_n. \end{aligned}$$

We have the picture of Σ shown in Fig. 8. In Fig. 8 we are using (4.4) which tells us that, since $d_r = 0$, (h_{k-1}) and (h_{k+1}) are basic special segments of type I. Since $a(h) = 8$ and $[u, v]$ is oriented toward v by h , we then have the picture of $\tilde{\Sigma}$ shown in Fig. 9. Figure 9 shows that $\tilde{\Sigma}$ is a simplex over \tilde{J} , relative to the partition $(\tilde{x}_0 < \tilde{y}_1 \leq \cdots \leq \tilde{y}_n < \tilde{x}_{n+1})$ of \tilde{J} , and that $\tilde{\mathbf{h}}$ and $\tilde{\Sigma}$ are engaged.

We note that (4.4) says that $r \in \mathcal{S}_2$, so there is a basic extraspecial segment \mathbf{x} of \mathbf{h} , such that in the notation of (4.2), we have

$$(h_{r-1}) = \mathbf{s}_{i-1}, \quad h_r = h_{k_1}, \quad (h_{r+1}) = \mathbf{s}_i,$$




FIGURE 8

for some i with $1 < i \leq u$. Put $\tilde{\mathbf{s}} = (h_{r-1}^h, h, h_{r+1})$ and put

$$\tilde{\mathbf{x}} = ((h_{k_1}) \circ \mathbf{s}_1 \circ \cdots \circ (h_{k_{i-1}}))^h \circ \tilde{\mathbf{s}} \circ (h_{k_{i+1}}) \circ \cdots \circ \mathbf{s}_{k_u} \circ (h_{k_{u+1}}).$$

One sees from the picture of $\tilde{\Sigma}$ that $\tilde{\mathbf{s}}$ is special of type I, relative to $\tilde{\Sigma}$. (See (3.6) for this.) Also (4.4) says that $m_k = 3$, so (3.9)(b) applied to \mathbf{x} yields $m_{k_{i-1}} \geq 6$ and $m_{k_{i+1}} \geq 6$. Thus, $\tilde{\mathbf{x}}$ satisfies (3.9)(b). The other parts of (3.9) pass unchanged from \mathbf{x} to $\tilde{\mathbf{x}}$, so $\tilde{\mathbf{x}}$ is extraspecial relative to $\tilde{\Sigma}$. Take $\mathbf{x} = \mathbf{x}_k$ in the decomposition (4.1). We then get

$$\tilde{\mathbf{h}} = (\mathbf{a}_1 \circ \mathbf{x}_1 \circ \cdots \circ \mathbf{x}_{k-1} \circ \mathbf{a}_k)^h \circ \tilde{\mathbf{x}} \circ \mathbf{a}_{k+1} \circ \cdots \circ \mathbf{x}_m \circ \mathbf{a}_{m+1},$$

and this decomposition shows that $\tilde{\mathbf{h}}$ and $\tilde{\Sigma}$ are married. This proves (5.6).

5.7. LEMMA. Suppose $\mathcal{S}_4 = \{r, r+1\}$, $y_r \neq y_{r+1}$, $a(h) = 12$, and $d(u, y_r) = d(y_r, y_{r+1}) = d(y_{r+1}, v) = 2$. Then $\tilde{J} \in \mathcal{P}(\mathbf{h}' \circ (h))$.

Proof. Define the points \tilde{x}_i , $0 \leq i \leq n+1$, the simplex $\tilde{\Sigma}$, and the points \tilde{y}_j , $1 \leq j \leq n$, exactly as in the proof of (5.6).

By (4.9), we have two cases to consider here, one in which $r \in \mathcal{S}_0$ and $m_r = 2$, and the other where $r \in \mathcal{S}_2$ and $m_r = 6$. In the second of these cases we get further information from (4.4)(iii). In particular, $d_{k-1} = d_{k+1} = 4$ if $r \in \mathcal{S}_2$. If $r \in \mathcal{S}_0$ then since $m_r = 2$ and $m(h) = 3$, (4.8) yields




FIGURE 9




FIGURE 10

$m_{r-1} \geq 6$ and $m_{r+1} \geq 6$. Here $\{r-1, r+1\} \subseteq \mathcal{S}_0 \cup \mathcal{S}_1$, so it follows from (3.9)(c) and (3.10) that $d_{r-1} \geq 4$ and $d_{r+1} \geq 4$. Thus, in any case we have the picture of σ shown in Fig. 10.

Since $a(h) = 12$, this yields the picture of $\tilde{\Sigma}$ shown in Fig. 11.

Suppose first that $r \in \mathcal{S}_0$. This leads to the consideration of four cases, according to whether $r \pm 1 \in \mathcal{S}_0$, $r-1 \in \mathcal{S}_0$ and $r+1 \in \mathcal{S}_1$, $r-1 \in \mathcal{S}_1$, and $r+1 \in \mathcal{S}_0$, and the case $r \pm 1 \in \mathcal{S}_1$. In any case, the above picture shows that (h) is a special segment of $\tilde{\mathbf{h}}$, of type II, relative to $\tilde{\Sigma}$. Write $\mathbf{h} = \mathbf{y} \circ \mathbf{b} \circ (h_r) \circ \mathbf{c} \circ \mathbf{z}$, where \mathbf{b} is a basic extraspecial segment if $r-1 \in \mathcal{S}_1$, and $\mathbf{b} = (h_{r-1})$ if $r-1 \in \mathcal{S}_0$, and where \mathbf{c} is defined similarly, relative to $r+1$. Put $\tilde{\mathbf{x}} = \mathbf{b}^h \circ (h) \circ \mathbf{c}$.

We claim that $\tilde{\mathbf{x}}$ is an extraspecial segment of $\tilde{\mathbf{h}}$, relative to $\tilde{\Sigma}$. Since $m_{r \pm 1} \geq 6$, as we have seen, we need only verify (3.9)(c) for $\tilde{\mathbf{x}}$. By symmetry, it will suffice to show

$$l(\tilde{\Sigma} \cap A(h_{r+1})) \geq 2 \cdot m_{r+1} - p - q((h)),$$

where $p = 0$ if $r+1 \in \mathcal{S}_0$, where $p = p(\mathbf{s})$ if $r+1 \in \mathcal{S}_1$ and \mathbf{s} is the basic special segment following (h_{r+1}) , and that the above inequality is strict if $\partial(\tilde{\Sigma} \cap A(h_{r+1})) \subseteq D$. The argument for this is perhaps familiar from the proof of (5.6), but we will supply it again.

We have $l(\tilde{\Sigma} \cap A(h_{r+1})) = d(y_{r+1}, x_{r+2}) = d(x_{r+1}, x_{r+2}) - d(x_{r+1}, y_{r+1})$. Here (3.9)(c) and (3.10) yield

$$(x_{r+1}, x_{r+2}) \geq 2 \cdot m_{r+1} - p,$$




FIGURE 11

with strict inequality if $\{x_{r+1}, x_{r+2}\} \subseteq D$, and (3.5) yields $d(x_{r+1}, y_{r+1}) \leq 2$, with strict inequality unless $\{x_{r+1}, y_{r+1}\} \subseteq D$. Now simply observe that $q((h)) = 2$ since $m(h) = 3$, in accordance with (3.8). This yields

$$d(y_{r+1}, x_{r+2}) \geq 2 \cdot m_{r+1} - p - q((h))$$

with the required condition concerning strict inequality, as desired. Thus, $\tilde{\mathbf{x}}$ is extraspecial relative to $\tilde{\Sigma}$. As in the proofs of (5.5) and (5.6), it now follows that $\tilde{\mathbf{h}}$ and $\tilde{\Sigma}$ are married. We are therefore finished with the case $r \in \mathcal{S}_0$.

We now have the case $r \in \mathcal{S}_2$ and $m_r = 6$. Let \mathbf{x} be the basic extraspecial segment of \mathbf{h} involving (h_r) . Then let \mathbf{s} and \mathbf{t} be the basic special segments preceding and, respectively, following (h_r) in \mathbf{x} . Put $\tilde{\mathbf{s}} = \mathbf{s}^h \circ (h) \circ \mathbf{t}$. By (4.4), \mathbf{s} and \mathbf{t} are of type II and $m_{k-1} = m_{k+1} = 2$. Then (3.6) shows that $l(\mathbf{s}) \equiv 2$ or $3 \pmod{4}$, and similarly for $l(\mathbf{t})$.

We claim that $\tilde{\mathbf{s}}$ is a special segment of $\tilde{\mathbf{h}}$, of type II, relative to $\tilde{\Sigma}$. Since $m(h) = 3$, $m(\tilde{\mathbf{s}})$ is indeed an alternating string of 2's and 3's, and it follows from our picture of $\tilde{\Sigma}$ that $\tilde{\mathbf{s}}$ satisfies the condition $(*)$ preceding (3.6). Our picture shows that, with respect to $\tilde{\Sigma}$, we have $d(h_{r-1}^h) = d_{r-1} - 2$ and $d(h_{r+1}) = d_{r+1} - 2$, and for any other entries in \mathbf{s} or \mathbf{t} , the value of the “ d -function” is unchanged on passage from Σ to $\tilde{\Sigma}$. Further, $d(h) = 6$ relative to $\tilde{\Sigma}$.

Suppose $l(\mathbf{s}) = l(\mathbf{t}) = 2$. Then $m(\tilde{\mathbf{s}}) = (3, 2, 3, 2, 3)$ and $d(\tilde{\mathbf{s}}) = (4, 2) \circ (6) \circ (2, 4)$, as required for (3.6)(II)(iv). Thus, $\tilde{\mathbf{s}}$ is special in this case. If $l(\mathbf{s}) = 2$ and $l(\mathbf{t}) = 3$ then $m(\tilde{\mathbf{s}}) = (3, 2, 3, 2, 3, 2)$ and $d(\tilde{\mathbf{s}}) = (4, 2) \circ (6) \circ (2, 2, 4)$, and again $\tilde{\mathbf{s}}$ is special. Now suppose $l(\mathbf{s}) = 2$ and $l(\mathbf{t}) > 4$. We then have $d(\mathbf{t}) = (4, 2, 2) \circ \xi \circ \mu$ where $\xi = \xi_t$, $t = [l(\mathbf{t})/4]$, and where $\mu = (2, 4)$ if $m(\mathbf{t})$ ends in 3, and $\mu = (2, 2, 4)$ if $m(\mathbf{t})$ ends in 2. From this we get

$$d(\tilde{\mathbf{s}}) = (4, 2) \circ (6) \circ (2, 2, 2) \circ \xi \circ \mu.$$

Since $l(\mathbf{t}) \not\equiv 1 \pmod{4}$, $[l(\tilde{\mathbf{s}})/4] = t + 1$, and $d(\tilde{\mathbf{s}}) = (4, 2) \circ \xi_{t+1} \circ \mu$, as required. The reader may now check that we also get $\tilde{\mathbf{s}}$ special of type II if $l(\mathbf{s}) = 3$. By symmetry, we may then assume that $l(\mathbf{s}) > 4$ and $l(\mathbf{t}) > 4$. Here we have $d(\mathbf{s}) = \lambda \circ \xi_s \circ (2, 2, 4)$, where $\lambda = (4, 2)$ if $m(\mathbf{s})$ begins with 3, and $\lambda = (4, 2, 2)$ if $m(\mathbf{s})$ begins with 2. Also, we write $d(\mathbf{t}) = (4, 2, 2) \circ \xi_t \circ \mu$, as above. Then

$$\begin{aligned} d(\tilde{\mathbf{s}}) &= \lambda \circ \xi_s \circ (2, 2, 2, 6, 2, 2, 2) \circ \xi_t \circ \mu \\ &= \lambda \circ \xi_{s+t+1} \circ \mu, \end{aligned}$$

and $s + t + 1 = [l(\tilde{\mathbf{s}})/4]$ since both $l(\mathbf{s})$ and $l(\mathbf{t})$ are congruent to 2 or 3, mod 4. Thus, $\tilde{\mathbf{s}}$ is special, of type II, in any case.

Let \mathbf{x} be the basic extraspecial segment of \mathbf{h} involving (h_r) , and in the notation of (4.2) put $\mathbf{s} = \mathbf{s}_{i-1}$, $\mathbf{t} = \mathbf{s}_i$. Then put

$$\tilde{\mathbf{x}} = \left((h_{k_1}) \circ \mathbf{s}_1 \circ \cdots \circ (h_{k_{i-1}}) \right)^h \circ \tilde{\mathbf{s}} \circ (h_{k_{i+1}}) \circ \cdots \circ \mathbf{s}_{k_u} \circ (h_{k_{u+1}}).$$

It now follows that $\tilde{\mathbf{x}}$ is an extraspecial segment of $\tilde{\mathbf{h}}$ relative to $\tilde{\Sigma}$. Taking $\mathbf{x} = \mathbf{x}_k$ is the notation of (4.1), we have

$$\tilde{\mathbf{h}} = (\mathbf{a}_1 \circ \mathbf{x}_1 \circ \cdots \circ \mathbf{a}_k)^h \circ \tilde{\mathbf{x}} \circ \mathbf{a}_{k+1} \circ \cdots \circ \mathbf{x}_m \circ \mathbf{a}_{m+1},$$

and evidently $\tilde{\mathbf{h}}$ and $\tilde{\Sigma}$ are married. This completes the proof of (5.7).

5.8. THEOREM. \mathcal{P} controls convergence relative to (b, \mathcal{D}) .

Proof. This is simply the union of the results (5.1) through (5.7), with help from (4.9).

5.9. COROLLARY. \mathcal{H} has the (b, \mathcal{D}) -convergence property.

Proof. This is immediate from (5.8) and from the main result of [5].

6. THE MAIN RESULTS

Recall that we have $F = F(S)$, the free group with free basis S . For any subset T of S , identify $F(T)$ with the subgroup of F generated by T . Then let X_T denote the subtree of X induced on the vertex-set $\{g, \langle a \rangle g : g \in F(T), a \in T\}$. Put $N = \langle \mathcal{H} \rangle$. Recall $N_{a,b} = \langle h_{a,b}^{F(\{a,b\})} \rangle$.

6.1. THEOREM. Let T be a subset of S , and put $N_T = \langle N_{a,b}^g : g \in F(T), a, b \in T, a \neq b \rangle$. Then $N \cap F(T) = \langle \mathcal{H} \cap F(T) \rangle = N_T$.

Proof. Evidently $N \cap F(T) \supseteq \langle \mathcal{H} \cap F(T) \rangle \supseteq N_T$. Assume (6.1) to be false and let $x \in N \cap F(T)$ with $x \notin N_T$. Among all such x , let x be chosen so that the length $n+1$ of x as a word in the alphabet \mathcal{H} is as small as possible. Let $\mathbf{h} = (h_0, \dots, h_n)$ be such a minimal-length word for x . Then $\mathbf{h} \in \mathcal{H}^*$ by (2.0)(a), and by Corollary (5.9) we may assume that x converges to 1 via \mathbf{h} , relative to (b, \mathcal{D}) . In particular, the closed interval $[x, 1] \cap A(h_0)$ has length at least $\frac{1}{2}a(h_0)$. But $a(h_0) \geq b(h_0) = 4 \cdot m(h_0) \geq 8$. This certainly implies that $[x, 1] \cap A(h)$ contains a segment I of the form $[\langle a \rangle g, \langle b \rangle g]$ for some $a, b \in S$ with $a \neq b$, and some $g \in F$. But $[x, 1] \subseteq X_T$, so $g \in F(T)$ and $a, b \in T$. Now (1.7)(d) and (1.6) together yield $h_0 \in N_{a,b}^g \leq N_T$.

Now $xh_0 \in N \cap F(T)$ and $xh_0 = (h_1 \cdots h_n)^{-1}$, so the minimality of the length of x yields $xh_0 \in N_T$, and then $x \in N_T$, a contradiction.

6.2. COROLLARY. *Let T be a subset of S , and let M_T be the Coxeter matrix $(m_{a,b})$ for $a, b \in T$. Then the canonical projection of $F(T)$ onto the Artin group $G(M_T)$ has the kernel $N \cap F(T)$. Thus, $G(M_T)$ may be identified with the subgroup $F(T)N/N$ of the Artin group G .*

Proof. The kernel of the canonical projection of $F(T)$ onto $G(M_T)$ is precisely the group N_T defined in (6.1). Then (6.1) yields a canonical isomorphism $F(T)N/N \cong G(M_T)$.

Corollary (6.2) is simply a re-statement of Theorem 2. It now remains to take up the Word Problem and Theorem 1. For the remainder of this paper, we assume that S is finite.

Put $S' = S \cup S^{-1}$, and let $FM(S')$ denote the free monoid on S' . Thus, elements of $FM(S')$ are finite sequences of elements of S' , and they will be referred to as *words*. The product $w \circ w'$ of two words is obtained by the usual operation of concatenating sequences. A word $w = (x_1, \dots, x_l)$ is *reduced* if $x_{i+1} \neq x_i^{-1}$ for any i , $1 \leq i < l$. The set of reduced words will be denoted by $\text{Red}(S')$. There is a natural bijection $\beta: \text{Red}(S') \rightarrow F$ given by $\beta(w) = x_1 \cdots x_l$.

A *section* of a word w is defined to be a triple (w_1, v, w_2) where $w = w_1 \circ v \circ w_2$.

Using β , the notions of *length* and of *syllable-length* lift from F to $\text{Red}(S')$. Thus, $|w|$ is simply the length of w as a sequence, while

$$\|w\| = \|\beta(w)\| = m$$

if we can write

$$g = \beta(w) = a_{i_1}^{e_1} \cdots a_{i_m}^{e_m}$$

as in (1.2). A section (w_1, v, w_2) of w will be said to be *syllable-complete* if we have

$$\beta(w_1) = a_{i_1}^{e_1} \cdots a_{i_{j-1}}^{e_{j-1}}, \quad \beta(v) = a_{i_j}^{e_j} \cdots a_{i_k}^{e_k}, \quad \beta(w_2) = a_{i_{k+1}}^{e_{k+1}} \cdots a_{i_m}^{e_m}$$

for some indices j and k , with $1 \leq j \leq k \leq m$. Thus, (w_1, v, w_2) is syllable-complete if w_1 doesn't end in the same letter with which v begins, and w_2 doesn't begin with the same letter in which v ends. Let $\mathcal{S}(w)$ denote the set of all syllable-complete sections of w . For any $a, b \in S$ with $a \neq b$, put

$$\mathcal{S}_{a,b}(w) = \{(w_1, v, w_2) \in \mathcal{S}(w) : \beta(v) \in \langle a, b \rangle\},$$

$$\mathcal{S}_{a,b}^*(w) = \{(w_1, v, w_2) \in \mathcal{S}_{a,b}(w) : \|v\| = m_{a,b}\}.$$

We then set

$$\begin{aligned}\mathcal{S}_{a,b} &= \cup\{\mathcal{S}_{a,b}(w) : w \in \text{Red}(S')\}, \\ \mathcal{S}_{a,b}^* &= \cup\{\mathcal{S}_{a,b}^*(w) : w \in \text{Red}(S')\}.\end{aligned}$$

Let $G_{a,b}$ denote the Artin group with generators a and b , and with the single relation $h_{a,b} = 1$ (from (0.2)). Identify $G_{a,b}$ with $\langle a, b \rangle / N_{a,b}$, where $N_{a,b}$ is the smallest normal subgroup of $\langle a, b \rangle$ containing $h_{a,b}$.

6.3. DEFINITION. Let $w \in \text{Red}(S')$ and let $a, b \in S$ with $a \neq b$. Define three subsets of $\mathcal{S}_{a,b}$, as follows.

$\tilde{\mathcal{M}}_{a,b}(w)$ = the set of all $(w_1, u, w_2) \in \mathcal{S}_{a,b}$ such that $\|u\| \leq m_{a,b}$, and such that there exists $v \in \text{Red}(S')$ with $(w_1, v, w_2) \in \mathcal{S}_{a,b}(w)$ and with $\beta(u^{-1}v) \in N_{a,b}$.

$\mathcal{M}_{a,b}(w)$ = the set of all $(w_1, u, w_2) \in \tilde{\mathcal{M}}_{a,b}(w)$ such that either $|u| < |v|$, or else $|u| = |v|$ and $\|u\| < \|v\|$, where $(w_1, v, w_2) \in \mathcal{S}_{a,b}(w)$.

$\mathcal{M}_{a,b}^*(w)$ = the set of all $(w_1, u, w_2) \in \tilde{\mathcal{M}}_{a,b}(w)$ such that $|u| = |v|$ and $\|u\| = \|v\| = m_{a,b}$, where $(w_1, v, w_2) \in \mathcal{S}_{a,b}(w)$.

6.4. PROPOSITION. For any $w \in \text{Red}(S')$, and for any $a, b \in S$ with $a \neq b$, the sets $\tilde{\mathcal{M}}_{a,b}(w)$, $\mathcal{M}_{a,b}(w)$, and $\mathcal{M}_{a,b}^*(w)$ are effectively constructible. Further, we have

$$\tilde{\mathcal{M}}_{a,b}(w) = \mathcal{M}_{a,b}(w) \sqcup \mathcal{M}_{a,b}^*(w) \sqcup \{(w_1, v, w_2) \in \mathcal{S}_{a,b}(w) : \|v\| < m_{a,b}\}.$$

Proof. The set $\mathcal{S}(w)$ of syllable-complete sections of w is constructible in quadratic time in $|w|$. For any $(w_1, v, w_2) \in \mathcal{S}_{a,b}(w)$, and any $u \in \text{Red}(\{a, b, a^{-1}, b^{-1}\})$ with $\|u\| \leq m_{a,b}$, (1.4)(a) says that if $\beta^{-1}(uv) \in N_{a,b}$ then $|u| \leq |v|$. Thus, in order to construct $\tilde{\mathcal{M}}_{a,b}(w)$, we may proceed as follows. First, construct $\mathcal{S}_{a,b}(w)$ (in quadratic time). Then, for each $(w_1, v, w_2) \in \mathcal{S}_{a,b}(w)$, make a list of all $u \in \text{Red}(\{a, b, a^{-1}, b^{-1}\})$ with $|u| \leq |v|$, and with $\|u\| \leq m_{a,b}$. (There are on the order of at most $4^{|w|}$ such words u .) For any such u , one can decide, in polynomial time on $|v|$, whether $\beta(u^{-1}v)$ is in $N_{a,b}$. This follows from the construction of canonical forms for elements of $G_{a,b}$, as given in [3, p. 677]. Among all such n for which $\beta(u^{-1}v) \in N_{a,b}$, choose only those u such that (w_1, u, w_2) is syllable-complete. In this way we have constructed $\tilde{\mathcal{M}}_{a,b}(w)$, in exponential time on $|w|$. The subsets $\mathcal{M}_{a,b}(w)$ and $\mathcal{M}_{a,b}^*(w)$ are then easily constructed, as well.

Now let $(w_1, u, w_2) \in \tilde{\mathcal{M}}_{a,b}(w)$, and let v be the element of $\text{Red}(S')$ with $(w_1, v, w_2) \in \mathcal{S}_{a,b}(w)$. As observed above, we have $|u| \leq |v|$, from (1.4)(a). Suppose that $(w_1, u, w_2) \notin \mathcal{M}_{a,b}(w)$. Then $|u| = |v|$ and $m_{a,b} \geq \|u\| \geq \|v\|$.

If now $\|v\| < m_{a,b}$, then (1.4)(b) yields $u = v$, and so $(w_1, v, w_2) \in \tilde{\mathcal{M}}_{a,b}(w)$. On the other hand, if $m_{a,b} = \|u\| = \|v\|$ then $(w_1, u, w_2) \in \mathcal{M}_{a,b}^*(w)$. This proves the second part of the proposition.

We can now state an algorithm which (as will be shown) solves the Word Problem for our locally non-spherical Artin group G .

6.5. ALGORITHM \mathcal{A} .

- (0) **Start** with a given $\tilde{w} \in \text{Red}(S')$. Put $\mathcal{L} = \{\tilde{w}\}$ and $\mathcal{L}' = \emptyset$. If $\tilde{w} = (\emptyset)$ go to (YES). Otherwise, go to (1).
 - (1) For every $w \in \mathcal{L}$ and every $a, b \in S$ with $a \neq b$, construct the sets $\mathcal{M}_{a,b}(w)$ and $\mathcal{M}_{a,b}^*(w)$. If $\mathcal{M}_{a,b}(w) \neq \emptyset$ for some w and some a and b , go to (2). Otherwise go to (2*).
 - (2) Choose $w \in \mathcal{L}$ and $a, b \in S$ with $\mathcal{M}_{a,b}(w) \neq \emptyset$, and then choose $(w_1, u, w_2) \in \mathcal{M}_{a,b}(w)$. If $w_1 \circ u \circ w_2 = (\emptyset)$, go to (YES). Otherwise replace \mathcal{L} by $\{w_1 \circ u \circ w_2\}$ and replace \mathcal{L}' by \emptyset , and go to (1).
 - (2*) Construct the set \mathcal{L}'' of all $w' = w_1 \circ u \circ w_2$ such that $(w_1, u, w_2) \in \mathcal{M}_{a,b}^*(w)$ for some $w \in \mathcal{L}$ and some $a, b \in S$. If $\mathcal{L}'' \subseteq \mathcal{L} \cup \mathcal{L}'$ go to (NO). Otherwise replace \mathcal{L}' with $\mathcal{L} \cup \mathcal{L}'$, and replace \mathcal{L} with $\mathcal{L}'' - (\mathcal{L} \cup \mathcal{L}')$, and go to (1).
- (YES) means that \tilde{w} represents 1 in G .
 (NO) means that \tilde{w} does not represent 1 in G .

The remainder of this section will be devoted to the proof that \mathcal{A} is an effective algorithm which solves the Word Problem for G . As a first step, it is important to see that \mathcal{A} does indeed return an answer of (YES) or (NO) to any $\tilde{w} \in \text{Red}(S')$.

Apart from the initial step from (0) to (1) (or to (YES)), and apart from the final step from (2) to (YES), or from (2*) to (NO), the algorithm \mathcal{A} consists of a sequence of loops, each of which is of the form (1) \rightarrow (2) \rightarrow (1) or (1) \rightarrow (2*) \rightarrow (1). A trivial induction argument then shows that at any stage, the list \mathcal{L} consists of words whose lengths are pair-wise equal. Now, in the course of a loop (1) \rightarrow (2) \rightarrow (1), \mathcal{L} is replaced by a “new” \mathcal{L} which consists of a singleton $\{w'\}$, and w' then has the property that $|w'| < |w|$ for any member w of any “old” version of \mathcal{L} . Thus, the “new” \mathcal{L} is disjoint from the union of all the old versions of \mathcal{L} in this case. On the other hand, in the course of a loop (1) \rightarrow (2*) \rightarrow (1), \mathcal{L} is replaced by a “new” \mathcal{L} which is disjoint from $\mathcal{L} \cup \mathcal{L}'$. But here $\mathcal{L} \cup \mathcal{L}'$ is the union of all the “old” versions of \mathcal{L} whose members are of the minimal length achieved by the algorithm so far. Since the members of the “new” \mathcal{L} also have this minimal length, it follows that, again, the “new” \mathcal{L} is disjoint from the union of all the old versions of \mathcal{L} .

Put $l = |\tilde{w}|$. There are then at most $\text{card}(S')^{(l^2+l)/2}$ reduced words of length at most l , so it now follows that after somewhat less than this

number of passages through (2) and (2*), the list \mathcal{L} will either be empty or will consist of the empty word. Thus, we have shown:

6.6. LEMMA. *For any $\tilde{w} \in \text{Red}(S')$, the algorithm \mathcal{A} returns an answer of (YES) or (NO), in exponential time relative to $|\tilde{w}|$.*

6.7. LEMMA. *Let $w \in \text{Red}(S')$, put $g = \beta(w)$, and consider the oriented closed segment $[g, 1]$ from g to 1 in X . Let $\mathcal{J}(w)$ denote the set of oriented closed segments $[g_1, g_2] \leq [g, 1]$, such that g_1 and g_2 are elements of F . There is a bijection θ_w from $\mathcal{J}(w)$ to the set $\mathcal{S}(w)$ of all syllable-complete sections of w , given by*

$$\theta_w([g_1, g_2]) = (\beta^{-1}(gg_1^{-1}), \beta^{-1}(g_1g_2^{-1}), \beta^{-1}(g_2)),$$

$$\theta_w^{-1}(w_1, v, w_2) = [\beta(v \circ w_2), \beta(w_2)].$$

Moreover, if $g_1g_2^{-1} \in \langle a, b \rangle$, where $a, b \in S$ and $a \neq b$, then $\theta_2([g_1, g_2]) \in \mathcal{S}_{a,b}(w)$.

Proof. For any $g' \in F$ we may write

$$g' = a_{i_1}^{e_1} \cdots a_{i_k}^{e_k},$$

where $a_{i_1}, \dots, a_{i_k} \in S$, $e_j \in \mathbb{Z}$, and $k = \|g'\|$. Put $x_0 = 1$, and define y_j and x_j , inductively, by

$$y_j = \langle a_{i_{k-j+1}} \rangle x_{j-1},$$

$$x_j = a_{i_{k-j+1}}^{e_{k-j+1}} x_{j-1},$$

for all j with $1 \leq j \leq k$. Then $g' = x_k$, and the vertex-sequence for the segment $[g, 1]$ is

$$(x_k, y_k, x_{k-1}, y_{k-1}, \dots, x_1, y_1, x_0).$$

Now take $g' = g$, where we are given $[g_1, g_2] \leq [g, 1]$. Thus, there exist indices r and s with $1 \leq r \leq s \leq k$, such that

$$g_1 = a_{i_r}^{e_r} \cdots a_{i_1}^{e_1}, \quad g_2 = a_{i_s}^{e_s} \cdots a_{i_1}^{e_1},$$

and it is then plain that $(\beta^{-1}(gg_1^{-1}), \beta^{-1}(g_1g_2^{-1}), \beta^{-1}(g_2))$ is a syllable-complete section of $w = \beta^{-1}(g)$, in $\mathcal{S}_{a,b}(w)$ if $g_1g_2^{-1} \in \langle a, b \rangle$.

On the other hand, let $(w_1, v, w_2) \in \mathcal{S}(w)$. We then have indices r and s as above, with

$$\beta(w_1) = a_{i_1}^{e_1} \cdots a_{i_{r-1}}^{e_{r-1}}, \quad \beta(v) = a_{i_r}^{e_r} \cdots a_{i_s}^{e_s}, \quad \beta(w_2) = a_{i_{s+1}}^{e_{s+1}} \cdots a_{i_k}^{e_k}.$$

It is then plain that $[\beta(vw_2), \beta(w_2)] \leq [g, 1]$, and that the maps that we have now obtained between $\mathcal{J}(w)$ and $\mathcal{S}(w)$ are inverse to each other.

6.8. Let $w \in \text{Red}(S')$ and put $g = \beta(w)$. We will denote by $\mathcal{H}(g)$ the set of all $h \in \mathcal{H}$ such that

$$d(gh, 1) + a(h) \leq d(g, 1) + b(h),$$

with strict inequality if $\partial(A(h) \cap [g, gh] \cap [g, 1]) \not\subseteq D$. (*)

For $h \in \mathcal{H}(g)$, let $I = I(g, h)$ denote the closed segment $A(h) \cap [g, gh] \cap [g, 1]$, oriented so that

$$I = [\partial_0 I, \partial_1 I] \leq [g, 1].$$

Notice that (0.4) and (*) together yield $l(I) \geq a(h) - \frac{1}{2}b(h)$. In particular, I is non-degenerate. We then define $\bar{I} = \bar{I}(g, h)$ to be the shortest, oriented closed segment such that

$$I \leq \bar{I} \leq [g, 1], \quad \text{with } \partial \bar{I} \subseteq D.$$

Thus, $l(\bar{I}) = l(I) + \text{card}(\partial I - D) \leq l(I) + 2$.

6.9. PROPOSITION. Let $w \in \text{Red}(S')$. Put $g = \beta(w)$, let $h \in \mathcal{H}(g)$, and put $w' = \beta^{-1}(gh)$. Then there exists $(w_1, u, w_2) \in \mathcal{M}_{a,b}(w) \cup \mathcal{M}_{a,b}^*(w)$ with $w_1 \circ u \circ w_2 = w'$.

Proof. Let g_1 and g_2 be the initial and terminal vertices, respectively, of $\bar{I}(g, h)$, so that $\bar{I}(g, h) = [g_1, g_2] \leq [g, 1]$. Let a and b be the (uniquely determined, by (1.7)(a)) elements of S such that $h \in \mathcal{H}_{a,b}$. Then $[g_1 h, g_2] \leq [gh, 1]$. Put $I = I(g, h)$ and $\bar{I} = \bar{I}(g, h)$. Also, put $J = A(h) \cap [g, gh] \cap [gh, 1]$, oriented so that $J \leq [gh, 1]$, and put $\bar{J} = [g_1 h, g_2]$. The situation is depicted in Fig. 12, which is drawn with J non-degenerate, and with $l(\bar{I}) = l(I) + 2$.

Put $(w_1, v, w_2) = \theta_2([g_1, g_2])$, as in (6.7), so that $v = \beta^{-1}(g_1 g_2^{-1})$. Put $h' = g_2 h g_2^{-1}$. Let a and b be the uniquely determined (by (1.7)(a)) elements of S such that $h \in \mathcal{H}_{a,b}$. Then (1.7)(c) shows that $h' \in N_{a,b}$ and $A(h') \subseteq X_{a,b}$. Since the vertex g_1 is at distance 1 from a vertex of $A(h)$ of the form $\langle a \rangle x$ or $\langle b \rangle x$, for some $x \in F$, it follows that $g_1 g_2^{-1}$ is at distance 1 from a vertex of $X_{a,b}$ of the form $\langle a \rangle x g_2^{-1}$ or $\langle b \rangle x g_2^{-1}$. But then $g_1 g_2^{-1}$ is a vertex of $X_{a,b}$, which is to say that $g_1 g_2^{-1} \in \langle a, b \rangle$. As $v = \beta^{-1}(g_1 g_2^{-1})$, we then have $(w_1, v, w_2) \in \mathcal{S}_{a,b}(w)$.

Put $u = \beta^{-1}(g_1 h g_2^{-1})$. Then $\beta(u^{-1}v) = (g_2 h^{-1} g_1^{-1})(g_1 g_2^{-1}) = (h')^{-1}$, so that $\beta(u^{-1}v) \in N_{a,b}$. Now, it should be clear from Fig. 12 (which comes, after all, by (0.4)) that $[g_1 h, g_2] \leq [gh, 1]$. Then (6.7) yields $\theta_{w'}([g_1 h, g_2]) = (w_1, u, w_2) \in \mathcal{S}_{a,b}(w')$. In order to obtain $(w_1, u, w_2) \in \tilde{\mathcal{M}}_{a,b}(w)$ it now remains to show that $\|u\| \leq m_{a,b}$.




FIGURE 12

The condition (6.8)(*) applies since $h \in \mathcal{H}(g)$, and so we have

$$a(h) - b(h) \leq l(I) - l(J).$$

Since $a(h) = l(I) + l(J)$, and since $b(h) = 4 \cdot m(h)$ by definition, we then obtain $l(J) \leq 2 \cdot m(h)$. Moreover, all of the inequalities are strict if $I \neq \bar{I}$. Notice also that if $l(\bar{J}) = l(J) + 2$ then $l(J)$ is even, and so $l(J) \leq 2 \cdot m(h) - 2$. We conclude that, in any case, we have $l(\bar{J}) \leq 2 \cdot m(h)$. Here $l(\bar{J}) = d(g_1 h, g_2) = d(\beta(u), 1)$, and so (1.3) gives $\|u\| \leq m(h)$. But $m(h) = m_{a,b}$ as $h \in \mathcal{H}_{a,b}$, so we obtain $(w_1, u, w_2) \in \tilde{\mathcal{M}}_{a,b}(w)$, as desired.

We have seen that $\beta(u^{-1}v) = (h')^{-1}$. Then $u \neq v$, as $h' \neq 1$. It now follows from the second statement in (6.4) that $(w_1, u, w_2) \in \mathcal{M}_{a,b}(w) \cup \mathcal{M}_{a,b}^*(w)$.

6.10. LEMMA. *Let $(w_1, u, w_2) \in \mathcal{M}_{a,b}^*(w)$. Put $w' = w_1 \circ u \circ w_2$, $g = \beta(w)$, $g' = \beta(w')$, and $h = g^{-1}g'$. Then $h \in \mathcal{H}(g) \cup \{1\}$.*

Proof. By assumption there exists $(w_1, v, w_2) \in \mathcal{S}_{a,b}^*(w)$ with $\beta(u^{-1}v) \in N_{a,b}$. If $u = v$ then $h = 1$ and we are done. So assume $u \neq v$ and put $h' = \beta(v^{-1}u)$. Then $h = g^{-1}g' = \beta(w^{-1})\beta(w') = \beta(w_2)^{-1}h'\beta(w_2)$. This shows that $h \in \mathcal{H}$. Now put $g_2 = \beta(w_2)$ and $g_1 = \beta(v \circ w_2)$. Then $[g_1, g_2] \leq [g, 1]$ by (6.7). Further, (6.7) shows that $d(g_1, g_2) = 2 \cdot \|v\|$ and $d(g_1 h, g_2) = 2 \cdot \|u\|$. Since $\|u\| = \|v\| = m_{a,b}$ we then get $d(g_1, g_1 h) \leq 4 \cdot m_{a,b}$. On the other hand, (1.12) yields $d(x, x \cdot h) \geq 4 \cdot m_{a,b}$ for any vertex x of X , with equality only if $a(h) = 4 \cdot m_{a,b}$, in which case (0.5) implies that $[x, x \cdot h]$ is a segment of $A(h)$. Thus, $[g_1, g_1 h]$ is a segment of $A(h)$ of length $a(h) = b(h)$. Now set $I = A(h) \cap [g, g'] \cap [g, 1]$. Then $[g_1, g_2] \leq I$,

and thus $l(I) \geq \frac{1}{2}a(h)$. It now follows that $d(g, 1) \geq d(g', 1)$. Moreover, the inequality is strict unless $I = [g_1, g_2]$. This shows that $h \in \mathcal{H}(g)$, in the sense of (6.8).

6.11. We may now prove that \mathcal{A} solves the Word Problem for G . Suppose false, and let $w \in \text{Red}(S')$. If \mathcal{A} returns an answer of (YES) to w , then it should be clear that w is indeed a word for the identity element of G . By (6.6) we may then assume that \mathcal{A} says (NO) to w , and that w is a word for the identity element. Among all such w , choose w so that $l = |w| + \|w\|$ is as small as possible.

Put $g = \beta(w)$. Then $g \in \langle \mathcal{H} \rangle$, and we may choose an \mathcal{H} -sequence $\mathbf{h} = (h_0, \dots, h_n)$ of minimal length for $g^{-1} = h_0 \cdots h_n$. By (2.0)(a) we then have $\mathbf{h} \in \mathcal{H}^*$ and $g^{-1} = h'_0 \cdots h'_n$ for any braiding (h'_0, \dots, h'_n) of \mathbf{h} . Then by Corollary (5.6) we may assume that \mathbf{h} has been chosen so that the vertex g converges to 1 via \mathbf{h} , relative to (b, \mathcal{D}) , in the sense of (2.5).

We now begin to follow w through \mathcal{A} . We have $w \neq (\emptyset)$ (else \mathcal{A} immediately says (YES)), so we go to step (1), with $\mathcal{L} = \{w\}$ and $\mathcal{L}' = \emptyset$. If $\mathcal{M}_{a,b}(w) \neq \emptyset$ for some $a, b \in S$ we go to (2), replacing w by some w' with $\beta(w') \in \langle \mathcal{H} \rangle$ and with $|w'| + \|w'\| < l$. By minimality of l , \mathcal{A} says (YES) to w' , hence to w also, for a contradiction. We conclude that $\mathcal{M}_{a,b}(w) = \emptyset$ for all $a, b \in S$ with $a \neq b$, and we go to (2*) instead of (2).

The algorithm \mathcal{A} will now continue to loop through steps (1) and (2*), until it sends us to (NO) (i.e., until \mathcal{L} becomes \emptyset). If at any stage we go to (2), we will reach a contradiction as above, via the minimality of l . On the other hand, for any k with $1 \leq k \leq n+1$, the k th passage through (2*) will yield $\beta^{-1}(gh_0 \cdots h_{k-1}) \in \mathcal{L}$. In order to see this, let us proceed slowly through (2*) for the k th time, $k \geq 1$. For any j with $1 \leq j \leq k$ put $x_j = \beta^{-1}(gh_0 \cdots h_{j-1})$, and put $x_0 = w = \beta^{-1}(g)$. Now form the list \mathcal{L}'' of all $w' = w_1 \circ u \circ w_2$ such that $(w_1, u, w_2) \in \mathcal{M}_{a,b}^*(w)$ for some $a, b \in S$. By induction on k , x_{k-1} is a member of the list \mathcal{L} as we enter step (2*). Here we have $\beta(x_{k-1}) = gh_0 \cdots h_{k-2}$, and we have $h_{k-1} \in \mathcal{H}(gh_0 \cdots h_{k-2})$ in the sense of (6.8), since g converges to 1 via \mathbf{h} . Now (6.9) yields $x_k = w_1 \circ u \circ w_2$ for some $(w_1, u, w_2) \in \mathcal{M}_{a,b}^*(w)$, and we get $x_k \in \mathcal{L}''$.

We now claim that $x_k \notin \mathcal{L} \cup \mathcal{L}'$. (Here \mathcal{L} and \mathcal{L}' have their “old” values. We haven’t progressed far enough through (2*), yet, to update \mathcal{L} and \mathcal{L}' .) Suppose false. Now, any element of $\mathcal{L} \cup \mathcal{L}'$ is constructed from w in the course of $k-1$ passes through (2*). It then follows from (6.10) that there are elements h'_1, \dots, h'_{k-1} of \mathcal{H} such that $x_k = \beta^{-1}(gh'_1 \cdots h'_{k-1})$. Then $\beta(x_k) = gh_0 \cdots h_{k-1} = gh'_1 \cdots h'_{k-1}$, and $g^{-1} = h'_1 \cdots h'_{k-1}h_k \cdots h_n$, contrary to the minimality of $l(\mathbf{h})$ as a word for g^{-1} in the alphabet \mathcal{H} . Thus $x_k \notin \mathcal{L} \cup \mathcal{L}'$, as claimed. We now finish step (2*), and arrive at a “new” \mathcal{L} with $x_k \in \mathcal{L}$.

In this way we finally get $x_{n+1} \in \mathcal{L}$. But $x_{n+1} = \beta^{-1}(gh_0 \cdots h_n) = (\emptyset)$. Since $|x_0| = |x_1| = \cdots = |x_{n+1}|$, it follows that w as (\emptyset) to begin with, for a contradiction. Thus, \mathcal{A} solves the Word Problem for G .

REFERENCES

1. K. I. Appel, On Artin groups and Coxeter groups of large type, in "Contributions to Group Theory" (Appel, Ratcliffe, and Schupp, Eds.), Contemporary Mathematics, Vol. 33, Amer. Math. Soc., Providence, 1984.
2. I. I. Appel and P. E. Schupp, Artin groups and infinite Coxeter groups, *Invent. Math.* **72** (1983), 201–220.
3. R. Charney, Artin groups of finite type are biautomatic, *Math. Ann.* **292** (1992), 671–683.
4. A. Chermak, \mathbb{R} -trees, small cancellation, and convergence, *Trans. Amer. Math. Soc.*, in press.
5. A. Chermak, A convergence theorem for Λ -trees, *J. Algebra*, in press.
6. J.-P. Serre, "Trees," Springer-Verlag, New York, 1980.
7. J. Tits, Sur le groupe des automorphismes d'un arbre, in "Essays on Topology and Related Topics: Memoires dédiés à Georges de Rham" (Haefliger and Narasimhan, Eds.), Springer-Verlag, New York/Berlin, 1970.